

Active Recombinant Human CD276 protein, His/Avi-tagged, Biotinylated

Cat. No. CD276-056H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant Human CD276(Leu29-Thr461) protein, fused to His/Avi tag at the C-terminus, was expressed in HEK293 cells .
Species	Human
Source	HEK293
ProteinLength	Leu29-Thr461
Description	<p>CD276 is a member of the B7 family of immune proteins that provide signals for the regulation of immune responses (1 - 3). Other family members include B7-1, B7-2, B7-H1/PD-L1, B7-H2, and PD-L2. B7 family proteins are type I transmembrane immunoglobulin (Ig) superfamily members that contain extracellular Ig V-like and Ig C-like domains with a short cytoplasmic tail. Among the family members there is about 20 - 40% amino acid (aa) sequence identity. B7-H3 was initially reported to be a 316 aa type I transmembrane precursor protein that contained a signal sequence, an extracellular region with one V-type and one C-type Ig domain, a transmembrane segment and a short cytoplasmic tail (1). Subsequent studies have identified a second 110 kDa form whose precursor is 534 aa in length. Termed 4IgB7-H3 or B7-H3b, this molecule has two additional Ig-like domains (one V-type and one C-type) and shows a ubiquitous expression pattern (4, 5). It would appear that the human 4Ig form is the principal, if not the only form of B7-H3 (5). Its precursor contains a 26 aa signal sequence, a 435 aa extracellular</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

region, a 31 aa transmembrane domain, and a 42 aa cytoplasmic tail. The four Ig-like domains alternate between V-type and C-type, and apparently are the consequence of a V-C type tandem duplication (4, 5). B7-H3b is expressed on dendritic cells as well as activated T, B and NK cells (5). The mouse gene differs from that of human in that it cannot code for four Ig-like domains; only a V-type:C-type pair (4). Human B7-H3b binding to an undefined receptor has shown to be inhibitory to NK cells and cytokine release (6). It also seems to be required for late stage osteoblast differentiation (7).

Predicted N Terminal Leu29

Form Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.

Bio-activity Measured in a cell proliferation assay using mouse CD4+ cells in the presence of anti-CD3. The ED50 for this effect is 3-12 µg/mL. Measured by its binding ability in a functional ELISA. When Human CD276 Antibody is immobilized at 1.0 µg/mL, 100 µL/well, the concentration of Biotinylated Recombinant Human CD276 His-tag Avi-tag that produces 50% of the optimal binding response is approximately 3 -15 ng/mL.

Molecular Mass 70-90 kDa, under reducing conditions

Endotoxin <1.0 EU per 1 µg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Applications Bioactivity

Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

after reconstitution.3 months, -20 to -70 °C under sterile conditions after reconstitution.

Reconstitution Reconstitute at 200 µg/mL in PBS.

Conjugation Biotin

GENE INFORMATION

Gene Name [CD276](#) [CD276 molecule \[Homo sapiens \]](#)

Official Symbol [CD276](#)

Synonyms [CD276](#); [CD276 molecule](#); [CD276 antigen](#); [B7 H3](#); [B7H3](#); [B7RP 2](#); [B7 homolog 3](#); [costimulatory molecule](#); [B7-H3](#); [B7RP-2](#); [4lg-B7-H3](#);

Gene ID [80381](#)

mRNA Refseq [NM_001024736](#)

Protein Refseq [NP_001019907](#)

MIM [605715](#)


UniProt ID [Q5ZPR3](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



2 µg/lane Protein was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining.

Binding Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA