

Active Recombinant Human CD28 protein, Fc/Avi-tagged, Biotinylated

Cat. No. CD28-051H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant Human CD28(Asn19-Pro152) protein, fused to Fc/Avi tag at the C-terminus, was expressed in CHO cells .
Species	Human
Source	CHO
ProteinLength	Asn19-Pro152
Description	<p>CD28 is the founding member of a subfamily of structurally homologous costimulatory or immune checkpoint molecules of the immunoglobulin (Ig) superfamily. The CD28/B7 pathway has been shown to play a central role in immune responses against infection, autoimmune diseases, and graft rejection (1). CD28 subfamily members, including ICOS, CTLA4, PD1, PD1H, and BTLA, are characterized by a single Ig V-like extracellular domain (ECD), a transmembrane domain and an intracellular domain with highly conserved tyrosine-based signaling motifs (2). CD28 and ICOS predominately enhance T-cell activation, while CTLA-4, PD-1 and BTLA are inhibitory (1-3). The mature ECD of human CD28 shares 65% and 66% amino acid sequence identity with mouse and rat CD28, respectively. CD28 and CTLA-4, together with their ligands, B7-1 and B7-2, constitute one of the dominant co-stimulatory pathways that regulate T and B cell responses. CD28 is expressed on approximately 80% of CD4+ T cells and 50% of CD8+ T cells in humans, with the proportion of CD28-positive T cells declining with age (1-4). CD28 expression has</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

been identified on other cell lineages, including bone marrow stromal cells, plasma cells, neutrophils, and eosinophils (5). CD28 is expressed on the cell surface as a disulfide-linked homodimer and the covalent bond is required for maximum activity (6,7). Although both CTLA-4 and CD28 can bind to the same ligands, CTLA-4 binds to B7-1 and B7-2 with a 20-100 fold higher affinity than CD28. While, CD28 ligation is critical in promoting proliferation and effector function of conventional T cells, it also promotes the anti-inflammatory function of Treg cells. Thus, CD28 serves both pro- and anti-inflammatory roles depending on the cell type and context in which it is expressed (8). The CD28 pathway has effects both at amplifying signals initiated by the T cell receptor and on a variety of processes, including signaling, metabolism, transcription, epigenetic modifications, post-translational modifications, and RNA splicing (9). Our Avi-tag Biotinylated CD28 features biotinylation at a single site contained within the Avi-tag, a unique 15 amino acid peptide. Protein orientation will be uniform when bound to streptavidin-coated surface due to the precise control of biotinylation and the rest of the protein is unchanged so there is no interference in the protein's bioactivity.

Predicted N Terminal	Asn19
Form	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.
Bio-activity	Measured by its binding ability in a functional ELISA. When Recombinant Human CD80 Fc Chimera Protein is immobilized at 10 g/mL (100 µ/well), the concentration of Biotinylated Recombinant CD28 Fc Chimera Avi-tag that produces a 50% optimal binding response is found to be 0.5-3 g/mL.
Molecular Mass	120-140 kDa, under non-reducing conditions
Endotoxin	<0.10 EU per 1 µg of the protein by the LAL method.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Applications	Bioactivity
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 2 weeks, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Reconstitution	Reconstitute at 200 µg/mL in sterile deionized water.
Conjugation	Biotin

GENE INFORMATION


Gene Name	CD28 CD28 molecule [Homo sapiens]
Official Symbol	CD28
Synonyms	CD28; CD28 molecule; CD28 antigen (Tp44); CD28 antigen; Tp44; MGC138290;
Gene ID	940
mRNA Refseq	NM_001243077
Protein Refseq	NP_001230006
MIM	186760
UniProt ID	P10747

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Binding Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA