

Recombinant Human CD36 cell lysate

Cat. No. CD36-2539HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human CD36 / SCARB3 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

CD36 CD36 molecule (thrombospondin receptor) [Homo sapiens]

Official Symbol

CD36

Synonyms

CD36; CD36 molecule (thrombospondin receptor); CD36 antigen (collagen type I receptor, thrombospondin receptor); platelet glycoprotein 4; FAT; GP3B; GP4; GPIV; SCARB3; GPIIIB; PAS IV; PAS-4 protein; glycoprotein IIIb; cluster determinant 36; fatty acid translocase; platelet glycoprotein IV; scavenger receptor class B, member 3; leukocyte differentiation antigen CD36; CHDS7; PASIV; BDPLT10;

Gene ID

948

mRNA Refseq

NM_000072

Protein Refseq

NP_000063

MIM

173510

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P16671
Chromosome Location	7q11.2
Pathway	Adaptive Immune System, organism-specific biosystem; Adipocytokine signaling pathway, organism-specific biosystem; Adipocytokine signaling pathway, conserved biosystem; Antigen processing-Cross presentation, organism-specific biosystem; Class I MHC mediated antigen processing and presentation, organism-specific biosystem; Cross-presentation of particulate exogenous antigens (phagosomes), organism-specific biosystem;
Function	Gram-negative bacterial cell surface binding; Gram-positive bacterial cell surface binding; high-density lipoprotein particle binding; lipid binding; lipoprotein particle binding; lipoteichoic acid receptor activity; low-density lipoprotein particle binding; low-density lipoprotein receptor activity; low-density lipoprotein receptor activity; receptor activity; thrombospondin receptor activity; transforming growth factor beta binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA