

Active Recombinant Human CD3E & CD3D protein, His & Flag-tagged, Biotinylated

Cat. No. CD3E & CD3D-375H **Lot. No.** (See product label)

SPECIFICATION

Product Overview Biotinylated Human CD3 epsilon & CD3 delta is expressed from human HEK293 cells. It contains AA Asp 23 - Asp 126 (CD3E) & Phe 22 - Ala 105 (CD3D) (Accession # NP_000724.1 (CD3E) & AAH39035 (CD3D)). It is the biotinylated form of Human CD3 epsilon & CD3 delta Protein. The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with biotins using standard chemical labeling method. A standard biotin reagent (13.5 angstroms) is used in this product. The biotin to protein ratio is 2.5-4.5 as determined by the HABA assay.

Species Human

Source HEK293

ProteinLength Asp 23 - Asp 126 (CD3E) & Phe 22 - Ala 105 (CD3D)


Form Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Bio-activity Immobilized Biotinylated Human CD3E&CD3D Heterodimer Protein, Fc,His Tag&Fc,Flag Tag, primary amine labeling at 1 µg/mL (100 µL/well)on streptavidin precoated (0.2 µg/well) plate, can bind Monoclonal Anti-Human CD3 Antibody, Mouse IgG2a (Clone: OKT3), premium grade with a linear range of 2-20 ng/mL.

Molecular Mass The protein has a calculated MW of 39.2 kDa (CD3E) and 37.5 kDa (CD3D) . The

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



	protein migrates as 45-55 kDa on a SDS-PAGE gel under reducing (R) condition due to glycosylation.
Endotoxin	Less than 1.0 EU per µg by the LAL method.
Purity	>95% as determined by SDS-PAGE. >90% as determined by SEC-MALS.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 centigrade to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	Biotin

GENE INFORMATION

Gene Name	CD3E CD3e molecule, epsilon (CD3-TCR complex) [Homo sapiens]
Official Symbol	CD3E
Synonyms	CD3E; CD3e molecule, epsilon (CD3-TCR complex); CD3e antigen, epsilon polypeptide (TiT3 complex); T-cell surface glycoprotein CD3 epsilon chain; CD3-epsilon; T-cell surface antigen T3/Leu-4 epsilon chain; T-cell antigen receptor complex, epsilon subunit of T3; T3E; TCRE; FLJ18683;



Gene ID	916
mRNA Refseq	NM_000733
Protein Refseq	NP_000724
UniProt ID	P07766

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA