

Active Recombinant Human CD40 Protein, Fc-His-tagged

Cat. No. CD40-31H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human CD40 Protein(NP_001241.1)(Met 1-Arg193) was expressed in HEK293, fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Species	Human
Source	HEK293
ProteinLength	1-193 aa
Form	Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5.
Bio-activity	Measured by its binding ability in a functional ELISA. Immobilized recombinant human CD40 at 2 µg/ml (100 µl/well) can bind biotinylated human CD40L with a linear range of 7.8-125 ng/ml.
Molecular Mass	The recombinant human CD40/Fc is a disulfide-linked homodimer. The reduced monomer consists of 421 amino acids and has a predicted molecular mass of 47.3 kDa. As a result of glycosylation, the apparent molecular mass of rhCD40/Fc monomer migrates with an apparent molecular mass of 55-60 kDa in SDS-PAGE under reducing conditions.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Purity	> 85 % as determined by SDS-PAGE

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name CD40 CD40 molecule, TNF receptor superfamily member 5 [Homo sapiens]

Official Symbol CD40

Synonyms CD40; CD40 molecule, TNF receptor superfamily member 5; TNFRSF5, tumor necrosis factor receptor superfamily, member 5; tumor necrosis factor receptor superfamily member 5; Bp50; p50; CD40L receptor; CD40 type II isoform; B cell-associated molecule; B cell surface antigen CD40; B-cell surface antigen CD40; CD40 antigen (TNF receptor superfamily member 5); tumor necrosis factor receptor superfamily, member 5; nerve growth factor receptor-related B-lymphocyte activation molecule; CDW40; TNFRSF5; MGC9013;

Gene ID 958

mRNA Refseq NM_001250

Protein Refseq NP_001241

MIM 109535


UniProt ID P25942

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA