

Active Recombinant Human CD47 Protein, hlgG/His-tagged

Cat. No. CD47-14H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human CD47 (19-141aa, 362aa), fused to hlgG-His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.
Species	Human
Source	Insect Cells
ProteinLength	19-141
Description	This gene encodes a membrane protein, which is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The encoded protein is also a receptor for the C-terminal cell binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This gene has broad tissue distribution, and is reduced in expression on Rh erythrocytes. Alternatively spliced transcript variants have been found for this gene.
Form	Liquid
Bio-activity	Measured by its binding ability in a functional ELISA with Human SIRP alpha/CD172a. The ED50 range ≤ 100 ng/mL.
Molecular Mass	40.9 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	QLLFNKTKSVEFTFCNDTVVIPCFVTNMEAQNTTEVYVKWKFKGRDIYTFDGALNKS TVPTDFSSAKIEVSQLLKGDASLKMDKSDAVSHTGNYTCEVTELTREGETIIELKYRV VSWFSPNE
Endotoxin	< 1 EU/μg of protein (determined by LAL method)
Purity	> 95% by SDS-PAGE
Applications	SDS-PAGE, Bioactivity
Storage	Can be stored at 2 to 8 centigrade for 1 week. For long term storage, aliquot and store at -20 to -80 centigrade. Avoid repeated freezing and thawing cycles.
Concentration	0.5 mg/mL (determined by absorbance at 280nm)
Storage Buffer	Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

GENE INFORMATION

Gene Name	CD47 CD47 molecule [Homo sapiens (human)]
Official Symbol	CD47
Synonyms	CD47; CD47 molecule; IAP; OA3; MER6; leukocyte surface antigen CD47; CD47 antigen (Rh-related antigen, integrin-associated signal transducer); CD47 glycoprotein; Rh-related antigen; antigen identified by monoclonal antibody 1D8; antigenic surface determinant protein OA3; integrin associated protein; integrin-associated signal transducer
Gene ID	961

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


mRNA Refseq [NM_001777](#)

Protein Refseq [NP_001768](#)

MIM [601028](#)

UniProt ID [Q08722](#)

SDS-PAGE




3 μ g by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Biological Activity



Human SIRP alpha/CD172a is coated at 2 $\mu\text{g}/\text{mL}$ (100 $\mu\text{L}/\text{well}$) can bind Human CD47. The ED50 range ≤ 100 ng/mL.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA