

## Recombinant Mouse Cd47 protein, His-tagged

Cat. No. CD47-3094M Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Mouse Cd47(Gln19-Pro158) fused with His tag at C-terminal was expressed in HEK293.
<b>Species</b>	Mouse
<b>Source</b>	HEK293
<b>ProteinLength</b>	Gln19-Pro158
<b>Description</b>	CD47 is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The protein is also a receptor for the C-terminal cell-binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This protein has broad tissue distribution, and is reduced in expression on Rh erythrocytes.
<b>Form</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
<b>AA Sequence</b>	QLLFSNVNSIEFTSCNETVVIPCIVRNVEAQSTEEMFVKWKLNKSIFYIDGNKNSTTT DQNFTS AKISVSDLINGIASLKMDKRDAMVGNVTCEVTELSREGKTVIELKNRRTAFNT DQGSACSYYYYEG GCKLVSWFSPVDHHHHHH
<b>Endotoxin</b>	Less than 0.1 ng/ug (1 IEU/ug) as determined by LAL test.
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Quality Control Test</b>	61.3 kDa
<b>Storage</b>	<p>Lyophilized protein should be stored at &lt; -20 centigrade, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7 centigrade for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at &lt; -20 centigrade for 3 months.</p>
<b>Reconstitution</b>	<p>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</p> <p>It is not recommended to reconstitute to a concentration less than 100 µg/ml.</p> <p>Dissolve the lyophilized protein in ddH<sub>2</sub>O.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>

## GENE INFORMATION

<b>Gene Name</b>	Cd47 CD47 antigen (Rh-related antigen, integrin-associated signal transducer) [ <i>Mus musculus</i> ]
<b>Official Symbol</b>	Cd47
<b>Synonyms</b>	CD47; CD47 antigen (Rh-related antigen, integrin-associated signal transducer); leukocyte surface antigen CD47; Rh-related antigen; integrin-associated protein; IAP; Itgp; AA407862; AI848868; AW108519; 9130415E20Rik; B430305P08Rik;
<b>Gene ID</b>	16423
<b>mRNA Refseq</b>	NM_010581
<b>Protein Refseq</b>	NP_034711
<b>UniProt ID</b>	Q61735

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Chromosome Location</b>	16; 16 B5
<b>Pathway</b>	Cell surface interactions at the vascular wall, organism-specific biosystem; Cell-Cell communication, organism-specific biosystem; ECM-receptor interaction, organism-specific biosystem; ECM-receptor interaction, conserved biosystem; Hemostasis, organism-specific biosystem; Signal regulatory protein (SIRP) family interactions, organism-specific biosystem;
<b>Function</b>	protein binding; thrombospondin receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA