

Active Recombinant Human CD48 protein, His/Avi-tagged, Biotinylated

Cat. No. CD48-051H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Biotinylated Recombinant Human CD48(Gln27-Ser220) protein, fused to His/Avi tag at the C-terminus, was expressed in CHO cells .

Species

Human

Source

CHO

ProteinLength

Gln27-Ser220

Description

CD48, also known as BLAST-1, BCM-1, and SLAMF2, is a 65 kDa GPI-linked protein in the CD2 family of immunoglobulin superfamily proteins (1-3). The human CD48 cDNA encodes a 243 amino acid (aa) precursor that includes a 26 aa signal sequence, a 194 aa mature protein containing two Ig-like C2-type domains, and a 23 aa C-terminal propeptide (4). A soluble form of CD48 has been detected in the serum of lymphoid leukemia and arthritis patients (5). Human CD48 shares approximately 50% aa sequence identity with mouse and rat CD48. It shares 20%-26% aa sequence identity with comparable regions of human 2B4, BLAME, CD2F-10, CD84, CD229, CRACC, NTB-A, and SLAM. CD48 is expressed on most lineage-committed hematopoietic cells but not on hematopoietic stem cells or multipotent hematopoietic progenitors (4, 6). Among dendritic cells (DC), CD48 is selectively expressed on circulating myeloid DC and resident bone marrow and thymus DC (7). CD2, 2B4, and heparan sulfate function as CD48 ligands (8-10). CD48 is competent to transduce signals and can also trigger signaling through CD2 or

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

2B4 (8, 11). CD48-CD2 interactions promote T cell activation and class switching to IgG2a in B cells (8, 12). High affinity CD48-2B4 interactions can either promote or inhibit NK cell and cytotoxic T cell (CTL) activation (7, 11, 13, 14). CD48-2B4 ligation does not directly trigger CTL activity but enhances signaling from the T cell receptor (13). CD48-2B4 mediated inhibition of NK cell activity is distinct from MHC I-restricted mechanisms (15). CD48 expressed on NK cells is co-activating, whereas CD48 expressed on other cell types inhibits NK cell activation (14). Both CD48 expressing and non-expressing cells can be targets of NK cell or CTL-mediated lysis (13, 16). Our Avi-tag Biotinylated human CD48 features biotinylation at a single site contained within the Avi-tag, a unique 15 amino acid peptide. Protein orientation will be uniform when bound to streptavidin-coated surface due to the precise control of biotinylation and the rest of the protein is unchanged so there is no interference in the protein's bioactivity.

Predicted N Terminal Gln27

Form Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.

Bio-activity Measured by its binding ability in a functional ELISA. When Recombinant Human CD244 Fc Chimera is immobilized at 1 µg/mL (100 µ/well), the concentration of Biotinylated Recombinant Human CD48 His-tag Avi-tag that produces 50% of the optimal binding response is approximately 0.3-1.8 µg/mL.

Molecular Mass 43-51 kDa, under reducing conditions

Endotoxin <0.10 EU per 1 µg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Applications	Bioactivity
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Reconstitution	Reconstitute at 250 µg/mL in PBS.
Conjugation	Biotin

GENE INFORMATION


Gene Name	CD48 CD48 molecule [Homo sapiens]
Official Symbol	CD48
Synonyms	CD48; CD48 molecule; BCM1, CD48 antigen (B cell membrane protein) , CD48 molecule; CD48 antigen; BLAST; hCD48; mCD48; SLAMF2; TCT.1; BCM1 surface antigen; leukocyte antigen MEM-102; B-lymphocyte activation marker BLAST-1; CD48 antigen (B-cell membrane protein); BCM1; BLAST1; MEM-102;
Gene ID	962
mRNA Refseq	NM_001256030
Protein Refseq	NP_001242959
MIM	109530
UniProt ID	P09326

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



2 µg/lane Protein was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining.

Binding Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA