

Active Recombinant Human CD70 protein, Fc-tagged, R-PE labeled

Cat. No. CD70-639P **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Recombinant Human CD70 protein, Fc-tagged was Conjugated with NH2-Reactive R-PE(R-Phycoerythrin).

Species

Human

Source

HEK293

ProteinLength

Gln 39-Pro 193

Form

Lyophilized from sterile 50 mM Tris, pH 8.0. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.

Bio-activity

Measured by its binding ability in a functional ELISA. Immobilized Human CD27 His at 2 µg/ml (100 µl/well) can bind Human CD70 hFc, the EC50 of Human CD70 hFc is 15-60 ng/mL.

Molecular Mass

The recombinant human CD70/Fc is a disulfide-linked homodimeric protein. The reduced monomer consists of 413 amino acids and predicts a molecular mass of 45.5 kDa. As a result of different glycosylation, the rh CD70/Fc migrates as several bands with the approximate molecular mass of 55-60, 110-120 and 160-170 kDa corresponding to the monomeric, dimeric and trimeric form respectively in SDS-PAGE under reducing conditions.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	< 1.0 EU per µg protein as determined by the LAL method.
Purity	> 90 % as determined by SDS-PAGE.
Storage	Samples are stable for up to twelve months from date of receipt at -20°C to -80°C Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	R-PE

GENE INFORMATION

Gene Name	CD70 CD70 molecule [Homo sapiens]
Official Symbol	CD70
Synonyms	CD70; CD70 molecule; CD27LG, TNFSF7, tumor necrosis factor (ligand) superfamily, member 7; CD70 antigen; CD27L; CD27-L; CD27 ligand; Ki-24 antigen; surface antigen CD70; tumor necrosis factor ligand superfamily member 7; tumor necrosis factor (ligand) superfamily, member 7; CD27LG; TNFSF7;
Gene ID	970
mRNA Refseq	NM_001252
Protein Refseq	NP_001243
MIM	602840

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



UniProt ID

P32970

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA