

Recombinant Human CD74 Protein, His-tagged

Cat. No. CD74-170H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human CD74 protein with His tag was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	296
Description	<p>The protein encoded by this gene associates with class II major histocompatibility complex (MHC) and is an important chaperone that regulates antigen presentation for immune response. It also serves as cell surface receptor for the cytokine macrophage migration inhibitory factor (MIF) which, when bound to the encoded protein, initiates survival pathways and cell proliferation. This protein also interacts with amyloid precursor protein (APP) and suppresses the production of amyloid beta (Abeta). Multiple alternatively spliced transcript variants encoding different isoforms have been identified.</p>
Form	Lyophilized
Molecular Mass	19.3 kDa
AA Sequence	<p>MHRRRSRSCREDQKPVMDQQRDLISNNEQLPMLGRRPGAPESKCSR GALYTGFSI LVTL LLAGQATTAYFLYQQQGR LDKLTVTSQNLQLENLRMKLPKPPKPVSKMRMAT PLLMQALPMGALPQGPMQNA TKYGNMTE DHVMHLLQNADPLKVYPPLKGSFPENL RHLKNTMETIDWKVFESWMHHWLLFEMSRHSLEQKPTDAPPKVLTKCQEEVSHIP A</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	VHPGSFRPKCDENGNLPLQCYGSIGYCWCVFPNGTEVPNTRSRGHHNCSESLEL EDPSSGLGVTKQDLGPVPM
Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.
GENE INFORMATION	
Gene Name	CD74 CD74 molecule, major histocompatibility complex, class II invariant chain [Homo sapiens (human)]
Official Symbol	CD74
Synonyms	CD74; CD74 molecule, major histocompatibility complex, class II invariant chain; CD74 antigen (invariant polypeptide of major histocompatibility complex, class II

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

antigen associated) , DHLAG; HLA class II histocompatibility antigen gamma chain; gamma chain of class II antigens; HLA DR gamma; Ia associated invariant chain; MHC HLA DR gamma chain; p33; HLA-DR-gamma; MHC HLA-DR gamma chain; Ia-associated invariant chain; HLA-DR antigens-associated invariant chain; CD74 antigen (invariant polypeptide of major histocompatibility complex, class II antigen-associated); II; DHLAG; HLADG; Ia-GAMMA; FLJ98970;

Gene ID [972](#)

mRNA Refseq [NM_001025158](#)

Protein Refseq [NP_001020329](#)

MIM [142790](#)

UniProt ID [P04233](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA