

## Active Recombinant Human CD80 protein, His-tagged

Cat. No. CD80-560H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human CD80(Val35-Asn242) fused with His tag at C-terminal was expressed in NS0.
<b>Species</b>	Human
<b>Source</b>	Mammalian Cells
<b>ProteinLength</b>	Val35-Asn242
<b>Predicted N Terminal</b>	Val35
<b>Form</b>	Lyophilized from a 0.2 µm filtered solution in PBS.
<b>Bio-activity</b>	Measured by its ability to induce IL-2 secretion by Jurkat human acute T cell leukemia cells. Freeman, G.J. et al. (1993) Science 262:909. The ED50 for this effect is typically 0.025-0.15 µg/mL in the presence of PHA.
<b>Molecular Mass</b>	Predicted Molecular Mass: 51.3 (monomer) kDa;SDS-PAGE: 70-90 kDa, reducing conditions
<b>Endotoxin</b>	<0.10 EU per 1 µg of the protein by the LAL method.
<b>Purity</b>	>95%, by SDS-PAGE under reducing conditions and visualized by silver stain.
<b>Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.12 months from

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

date of receipt, -20 to -70 centigrade as supplied.1 month, 2 to 8 centigrade under sterile conditions after reconstitution.3 months, -20 to -70 centigrade under sterile conditions after reconstitution.

**Reconstitution** Reconstitute at 100 µg/mL in sterile PBS.

**Shipping** The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

## GENE INFORMATION

**Gene Name** CD80 CD80 molecule [ Homo sapiens ]

**Official Symbol** CD80

**Synonyms**

CD80; CD80 molecule; CD28LG, CD28LG1, CD80 antigen (CD28 antigen ligand 1, B7 1 antigen) , CD80 molecule; T-lymphocyte activation antigen CD80; B lymphocyte activation antigen B7; B7 1; B7.1; activation B7-1 antigen; costimulatory factor CD80; CTLA-4 counter-receptor B7.1; B-lymphocyte activation antigen B7; costimulatory molecule variant IgV-CD80; CD80 antigen (CD28 antigen ligand 1, B7-1 antigen); B7; BB1; B7-1; LAB7; CD28LG; CD28LG1;

**Gene ID** 941

**mRNA Refseq** NM\_005191

**Protein Refseq** NP\_005182

**MIM** 112203

**UniProt ID** P33681

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Chromosome Location</b>	3q13.3-q21
<b>Pathway</b>	Adaptive Immune System, organism-specific biosystem; Allograft rejection, organism-specific biosystem; Allograft rejection, conserved biosystem; Autoimmune thyroid disease, organism-specific biosystem; Autoimmune thyroid disease, conserved biosystem; CD28 co-stimulation, organism-specific biosystem; CD28 dependent PI3K/Akt signaling, organism-specific biosystem;
<b>Function</b>	coreceptor activity; protein binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA