

Active Recombinant Human CD80, His-tagged, Biotinylated

Cat. No. CD80-581H Lot. No. (See product label)

SPECIFICATION

Product Overview The recombinant human CD80 ECD protein is expressed as a 220 amino acid protein consisting of Val35 - Asn242 region of CD80 (UniProt accession #P33681) and a C-terminal His-tag. It contains 8 potential sites for N-linked glycosylation.

Species Human

Source Human Cells

ProteinLength 35-242 a.a.

Form Supplied at 0.5 mg/ml in sterile PBS pH7.4 (carrier free & preservative free). The purified recombinant protein was labeled with Biotin (3-5 Biotin per molecule).

Bio-activity Binds human CD28, CTLA4 and PD-L1. Induces IL-2 secretion in human T cells.

Molecular Mass Calculated molecular mass 25.3 kDa; estimated by SDS-PAGE under reducing condition 45-55 kDa probably due to glycosylation.

AA Sequence
 VIHVTKEVKEVATLSCGHNVSVEELAQTRIYWQKEKMMVLTMMSGDMNIWPEYKNR
 TIFDITNNLSIVILALRP SDEGTYECVVLKYEKDAFKREHLAEVTL SVKADFP TSPISDF
 EIPTSNIRRIICSTSGGFPEPHLSWLENGEEL NAINTTVSQDPETELYAVSSKLD FNM
 TTNHSFMCLIKYGHLRVNQTFNWNTTKQEHPDNGSTGHHHHHHHHH

Endotoxin <0.1 eu per 1 µg of purified recombinant protein determined by the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Purity	>95% judged by SDS-PAGE under reducing condition
Storage	The product is shipped at 4°C. Upon receipt, centrifuge the product briefly before opening the vial. It is recommended to store small aliquots at the temperature below -20°C for long-term storage and the product is stable for 3 months. The undiluted protein can be stored at 4°C for no more than 2 weeks. Avoid repeated freeze-thaw cycles.
Conjugation	Biotin

GENE INFORMATION

Gene Name	CD80 CD80 molecule [Homo sapiens]
Official Symbol	CD80
Synonyms	CD80; CD80 molecule; CD28LG, CD28LG1, CD80 antigen (CD28 antigen ligand 1, B7 1 antigen) , CD80 molecule; T-lymphocyte activation antigen CD80; B lymphocyte activation antigen B7; B7 1; B7.1; activation B7-1 antigen; costimulatory factor CD80; CTLA-4 counter-receptor B7.1; B-lymphocyte activation antigen B7; costimulatory molecule variant IgV-CD80; CD80 antigen (CD28 antigen ligand 1, B7-1 antigen); B7; BB1; B7-1; LAB7; CD28LG; CD28LG1;
Gene ID	941
mRNA Refseq	NM_005191
Protein Refseq	NP_005182
MIM	112203

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA



UniProt ID	P33681
Chromosome Location	3q13.3-q21
Pathway	Adaptive Immune System, organism-specific biosystem; Allograft rejection, organism-specific biosystem; Allograft rejection, conserved biosystem; Autoimmune thyroid disease, organism-specific biosystem; Autoimmune thyroid disease, conserved biosystem; CD28 co-stimulation, organism-specific biosystem; CD28 dependent PI3K/Akt signaling, organism-specific biosystem;
Function	coreceptor activity; protein binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA