

Active Recombinant Human CD80 Protein, Fc & Avi-tagged, Biotinylated

Cat. No. CD80-583H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant Human CD80 Protein (Val 35 - Asn 242) (NP_005182.1), fused with a human IgG1 Fc tag at the C-terminus, followed by a Avi tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	Val 35 - Asn 242
Predicted N Terminal	Val 35
Form	Lyophilized from sterile 50 mM tris, 100 mM glycine, pH7.5, 10% trehalose.
Bio-activity	Measured by its binding ability in a functional ELISA. Immobilized Human CTLA-4, Fc Tag at 5µg/mL (100 µL/well) can bind Biotinylated Human B7-1, Fc Tag with a linear range of 0.6-9.8 ng/mL.
Molecular Mass	Recombinant Human CD80 Protein has a calculated MW of 52.3 kDa. As a result of glycosylation, The protein migrates as 65-95 kDa under reducing (R) condition and 130-160 kDa under non-reducing (NR) condition on SDS-PAGE gel.
Endotoxin	Less than 1.0 EU per µg by the LAL method.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	>95% as determined by SDS-PAGE.
Stability	Samples are stable for up to twelve months from date of receipt at -70°C.
Storage	Store it under sterile conditions at -20°C~-70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	Biotin

GENE INFORMATION

Gene Name	CD80 CD80 molecule [Homo sapiens]
Official Symbol	CD80
Synonyms	CD80; CD80 molecule; CD28LG, CD28LG1, CD80 antigen (CD28 antigen ligand 1, B7 1 antigen) , CD80 molecule; T-lymphocyte activation antigen CD80; B lymphocyte activation antigen B7; B7 1; B7.1; activation B7-1 antigen; costimulatory factor CD80; CTLA-4 counter-receptor B7.1; B-lymphocyte activation antigen B7; costimulatory molecule variant IgV-CD80; CD80 antigen (CD28 antigen ligand 1, B7-1 antigen); B7; BB1; B7-1; LAB7; CD28LG; CD28LG1;
Gene ID	941
mRNA Refseq	NM_005191
Protein Refseq	NP_005182

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	112203
UniProt ID	P33681
Chromosome Location	3q13.3-q21
Pathway	Adaptive Immune System, organism-specific biosystem; Allograft rejection, organism-specific biosystem; Allograft rejection, conserved biosystem; Autoimmune thyroid disease, organism-specific biosystem; Autoimmune thyroid disease, conserved biosystem; CD28 co-stimulation, organism-specific biosystem; CD28 dependent PI3K/Akt signaling, organism-specific biosystem;
Function	coreceptor activity; protein binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA