

## Recombinant Human CD86 Protein, Fc-tagged

**Cat. No.** CD86-152H    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human protein from the extracellular domain(Leu26-His245) of homo sapiens CD86 molecule(CD86), transcript variant 1(NM_175862), fused with the extracellular domain(Pro134-Leu281) of homo sapiens Fas ligand(FASLG), transcript variant 1(NM_000639) , with a Fc tag, was expressed in human cells.
<b>Species</b>	Human
<b>Source</b>	Human Cells
<b>ProteinLength</b>	26-245;134-281 a.a.
<b>Description</b>	This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.
<b>Form</b>	Purified protein formulated in a sterile solution of TBS buffer, pH7.221, without any preservatives.
<b>Molecular Mass</b>	70 kDa

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Endotoxin</b>	Endotoxin level is < 0.1 ng/g of protein (<1EU /g)
<b>Purity</b>	>90% by SDS-PAGE gel and Coomassie Blue staining
<b>Applications</b>	Antigens, Western, ELISA and other in vitro binding or in vivo functional assays, and protein-protein interaction studies.

## GENE INFORMATION


<b>Gene Name</b>	CD86 CD86 molecule [ Homo sapiens ]
<b>Official Symbol</b>	CD86
<b>Synonyms</b>	CD86; CD86 molecule; CD28LG2, CD86 antigen (CD28 antigen ligand 2, B7 2 antigen); T-lymphocyte activation antigen CD86; B lymphocyte antigen B7 2; B7 2; B7.2; BU63; FUN-1; CTLA-4 counter-receptor B7.2; B-lymphocyte activation antigen B7-2; CD86 antigen (CD28 antigen ligand 2, B7-2 antigen); B70; B7-2; LAB72; CD28LG2; MGC34413;
<b>Gene ID</b>	942
<b>mRNA Refseq</b>	NM_001206924
<b>Protein Refseq</b>	NP_001193853
<b>MIM</b>	601020
<b>UniProt ID</b>	P42081

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA