

Recombinant Human CD93 cell lysate

Cat. No. CD93-1824HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human CD93 / C1QR1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

CD93 CD93 molecule [Homo sapiens]

Official Symbol

CD93

Synonyms

CD93; CD93 molecule; C1QR1, CD93 antigen , complement component 1, q subcomponent, receptor 1 , matrix remodelling associated 4 , MXRA4; complement component C1q receptor; C1qR(P); C1qRP; CDw93; dJ737E23.1; ECSM3; C1qR; CD93 antigen; C1q receptor 1; C1q/MBL/SPA receptor; matrix-remodelling associated 4; matrix-remodeling-associated protein 4; complement component 1 q subcomponent receptor 1; complement component 1, q subcomponent, receptor 1; C1QR1; MXRA4;

Gene ID

22918

mRNA Refseq

NM_012072

Protein Refseq

NP_036204

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



MIM	120577
UniProt ID	Q9NPY3
Chromosome Location	20p11.21
Function	calcium ion binding; complement component C1q binding; protein binding; receptor activity; sugar binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA