

## Recombinant Human CDC23 293 Cell Lysate

**Cat. No.** CDC23-7667HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for cell division cycle 23 homolog ( <i>S. cerevisiae</i> ) (CDC23) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	CDC23 cell division cycle 23 homolog ( <i>S. cerevisiae</i> ) [ <i>Homo sapiens</i> ]
<b>Official Symbol</b>	CDC23
<b>Synonyms</b>	CDC23; cell division cycle 23 homolog ( <i>S. cerevisiae</i> ); CDC23 (cell division cycle 23, yeast, homolog); cell division cycle protein 23 homolog; ANAPC8; anaphase promoting complex subunit 8; APC8; CUT23; cyclosome subunit 8; anaphase-promoting complex subunit 8;
<b>Gene ID</b>	8697
<b>mRNA Refseq</b>	NM_004661
<b>Protein Refseq</b>	NP_004652
<b>MIM</b>	603462
<b>UniProt ID</b>	Q9UJX2
<b>Chromosome Location</b>	5q31
<b>Pathway</b>	APC/C complex, organism-specific biosystem; APC/C complex, conserved biosystem; APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Cyclin B, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem; APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem;

**Function**

binding; ubiquitin-protein ligase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA