

## Recombinant Human CDC26 293 Cell Lysate

**Cat. No.** CDC26-7663HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for cell division cycle 26 homolog ( <i>S. cerevisiae</i> ) (CDC26) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	CDC26 cell division cycle 26 homolog ( <i>S. cerevisiae</i> ) [ <i>Homo sapiens</i> ]
<b>Official Symbol</b>	CDC26
<b>Synonyms</b>	CDC26; cell division cycle 26 homolog ( <i>S. cerevisiae</i> ); C9orf17, cell division cycle 26 , chromosome 9 open reading frame 17; anaphase-promoting complex subunit CDC26; ANAPC12; anaphase promoting complex subunit 12; APC12; CDC26 subunit of anaphase promoting complex; anaphase-promoting complex subunit 12; cell division cycle protein 26 homolog; C9orf17;
<b>Gene ID</b>	<a href="#">246184</a>
<b>mRNA Refseq</b>	<a href="#">NM_139286</a>
<b>Protein Refseq</b>	<a href="#">NP_644815</a>
<b>MIM</b>	<a href="#">614533</a>
<b>UniProt ID</b>	<a href="#">Q8NHZ8</a>
<b>Chromosome Location</b>	9q32
<b>Pathway</b>	APC/C complex, organism-specific biosystem; APC/C complex, conserved biosystem; APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Cyclin B, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem;  
APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted  
proteins in late mitosis/early G1, organism-specific biosystem;

**Function**

protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA