

Native Chicken CII

Cat. No. CII-250C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Collagen type II (CII) protein, purified from chicken sternum, for the induction of arthritis in the Collagen-Induced Arthritis (CIA) model.
Species	Chicken
Source	Chicken Sternum
Description	<p>Type II collagen (CII) is a fibrillar collagen that is primarily located in cartilage tissue. The major molecular form of collagen in cartilage is type II collagen. Type II collagen is also located in the vitreous humor of the eye, the inner ear, the nose, and the intervertebral discs of the spine. The function of type II collagen is to provide tensile strength and give cartilage the ability to resist shearing forces. Mutations of the COL2A1 gene affect the synthesis of type II collagen and cause chondrodysplasias. This is characterized by abnormal cartilage that leads to bone and joint deformities. The degradation of collagen type II is an early indicator of osteoarthritis. Type II Collagen and Adjuvant Susceptibility to CIA is linked to MHC class II molecules and is dependent upon the species of type II collagen used for immunization. Various species of highly purified Type II Collagen are supplied lyophilized and in solution for use in the induction of arthritis in vivo</p>
Form	Lyophilized?
Molecular Mass	300 kDa
Purity	≥99% as determined by SDS-PAGE and visualized by Coomassie blue staining

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Applications

Induction of arthritis in vivo; Cell Culture; ELISA standard/plate coating

Storage

Collage dissolved in acetic acid is stable at 4 °C for 1 month. May be stored in dark at -20°C for long term storage but will gradually degrade under neutral conditions.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA