

Active Recombinant Human CLDN18.2 Full Length Transmembrane protein, His-tagged

Cat. No. CLDN18.2-1039H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human CLDN18.2 protein(P56856-2)(1-261aa), fused with C-terminal His tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	1-261aa
Form	<p>If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol.</p> <p>If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.</p>
Bio-activity	Measured by its binding ability in a functional ELISA. Immobilized human CLDN18.2 at 5 µg/ml can bind anti-CLDN18.2 recombinant Monoclonal Antibody, the the EC50 is 5.225-9.256 ng/ml.
Molecular Mass	29.1 kDa
AA Sequence	<p>MAVTACQQLGFVVSLLIGIAGIIAATCMDQWSTQDLYNNPVTAVFNYQGLWRSCVRE</p> <p>SSGFTECRGYFTLLGLPAMLQAVRALMIVGIVLGAIGLLVSIFALKCIRIGSMEDSAKA</p> <p>NMTLTSGIMFIVSGLCAIAGVSVFANMLVTNFWMSTANMYTGMGGMVQTVQTRYTF</p> <p>GAALFVGWVAGGLTLIGGVMMCIACRGLAPEETNYKAVSYHASGHSVAYKPGGFKA</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

STGFGSNTKNKKIYDGGARTEDEVQSYP SKHDYV

Endotoxin

Less than 1.0 EU/ug as determined by LAL method.


Storage

Store at -20°C/-80°C upon receipt, aliquoting is necessary for mutiple use. Avoid repeated freeze-thaw cycles.

Reconstitution

Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C.

SDS-PAGE



Activity

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA