

# Recombinant Human CLEC7A cell lysate

**Cat. No.** CLEC7A-001HCL    **Lot. No.** (See product label)

## SPECIFICATION

<b>Product Overview</b>	Human CLEC7A / Dectin-1 / CLECSF12 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.
<b>Species</b>	Human
<b>Source</b>	Human Cells
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

## GENE INFORMATION

**Gene Name**

[CLEC7A C-type lectin domain family 7, member A \[ Homo sapiens \]](#)

**Official Symbol**

CLEC7A

**Synonyms**

CLEC7A; C-type lectin domain family 7, member A; C type (calcium dependent, carbohydrate recognition domain) lectin, superfamily member 12 , CLECSF12; C-type lectin domain family 7 member A; dectin 1; hDectin 1; dectin-1; beta-glucan receptor; lectin-like receptor 1; DC-associated C-type lectin 1; C-type lectin superfamily member 12; dendritic cell-associated C-type lectin 1; dendritic cell-associated C-type lectin-1; C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 12; BGR; CANDF4; DECTIN1; CLECSF12;

**Gene ID**

[64581](#)

**mRNA Refseq**

[NM\\_022570](#)

**Protein Refseq**

[NP\\_072092](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



<b>MIM</b>	606264
<b>UniProt ID</b>	Q9BXN2
<b>Chromosome Location</b>	12p13.2-p12.3
<b>Pathway</b>	Phagosome, organism-specific biosystem; Phagosome, conserved biosystem; Tuberculosis, organism-specific biosystem; Tuberculosis, conserved biosystem;
<b>Function</b>	(1->3)-beta-D-glucan binding; MHC protein binding; binding; metal ion binding; opsonin binding; pattern recognition receptor activity; sugar binding; sugar binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA