

Recombinant Human CRK 293 Cell Lysate

Cat. No. CRK-7277HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for v-crk sarcoma virus CT10 oncogene homolog (avian) (CRK), transcript variant I is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CRK v-crk sarcoma virus CT10 oncogene homolog (avian) [Homo sapiens]
Official Symbol	CRK
Synonyms	CRK; v-crk sarcoma virus CT10 oncogene homolog (avian); v crk avian sarcoma virus CT10 oncogene homolog; adapter molecule crk; proto-oncogene c-Crk; p38; CRKII; FLJ38130;
Gene ID	1398
mRNA Refseq	NM_005206
Protein Refseq	NP_005197
MIM	164762
UniProt ID	P46108
Chromosome Location	17p13
Pathway	ARMS-mediated activation, organism-specific biosystem; Angiopoietin receptor Tie2-mediated signaling, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; Bacterial invasion of epithelial cells, organism-specific biosystem; Bacterial invasion of epithelial cells, conserved biosystem; CXCR4-mediated signaling events, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

SH2 domain binding; SH3/SH2 adaptor activity; ephrin receptor binding; protein binding; protein phosphorylated amino acid binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA