

## Recombinant Human CRYAB, His-tagged

Cat. No. CRYAB-11598H Lot. No. (See product label)

### SPECIFICATION

**Product Overview** Recombinant Human CRYAB protein, fused to His-tag, was expressed in E.coli and purified by Ni-sepharose.

**Species** Human

**Source** E.coli

**ProteinLength** 1-175a.a.

#### Description

Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Alpha crystallins are composed of two gene products: alpha-A and alpha-B, for acidic and basic, respectively. Alpha crystallins can be induced by heat shock and are members of the small heat shock protein (sHSP also known as the HSP20) family. They act as molecular chaperones although they do not renature proteins and release them in the fashion of a true chaperone; instead they hold them in large soluble aggregates. Post-translational modifications decrease the ability to chaperone. These heterogeneous

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

aggregates consist of 30-40 subunits; the alpha-A and alpha-B subunits have a 3:1 ratio, respectively. Two additional functions of alpha crystallins are an autokinase activity and participation in the intracellular architecture. Alpha-A and alpha-B gene products are differentially expressed; alpha-A is preferentially restricted to the lens and alpha-B is expressed widely in many tissues and organs. Elevated expression of alpha-B crystallin occurs in many neurological diseases; a missense mutation cosegregated in a family with a desmin-related myopathy.

<b>Source</b>	E.coli
<b>Species</b>	Human
<b>Tag</b>	His
<b>Storage</b>	The protein is stored in PBS buffer at -20°C. Avoid repeated freezing and thawing cycles.
<b>Storage Buffer</b>	1M PBS (58mM Na <sub>2</sub> HPO <sub>4</sub> , 17mM NaH <sub>2</sub> PO <sub>4</sub> , 68mM NaCl, pH8. ) added with 300mM Imidazole and 0.7% Sarcosyl, 15% glycerol.

## GENE INFORMATION

<b>Gene Name</b>	CRYAB crystallin, alpha B [ Homo sapiens ]
<b>Official Symbol</b>	CRYAB
<b>Synonyms</b>	CRYAB; crystallin, alpha B; CRYA2; alpha-crystallin B chain; HSPB5; heat shock protein beta-5; rosenthal fiber component; heat-shock 20 kD like-protein; renal carcinoma antigen NY-REN-27; CTPP2;
<b>Gene ID</b>	1410

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>mRNA Refseq</b>	NM_001885
<b>Protein Refseq</b>	NP_001876
<b>MIM</b>	123590
<b>UniProt ID</b>	P02511
<b>Chromosome Location</b>	11q22.3-q23.1
<b>Pathway</b>	Protein processing in endoplasmic reticulum, organism-specific biosystem; Protein processing in endoplasmic reticulum, conserved biosystem;
<b>Function</b>	protein binding; protein homodimerization activity; structural constituent of eye lens; unfolded protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA