

Recombinant Human CSF1, His-tagged

Cat. No. CSF1-96H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human M-CSF, fused to His-tag at N-terminus, was expressed as insoluble protein aggregate in <i>E.coli</i> and purified by conventional chromatography, after refolding of the isolated inclusion bodies in a renaturation buffer. MW = 20.7 kDa (179aa).
Species	Human
Source	<i>E.coli</i>
Description	Macrophage colony stimulating factor (M-CSF), as known as CSF-1, one of the hematopoietic growth factors that regulate the growth and differentiation of blood cells. This protein is produced by monocytes, granulocytes, endothelial cells, and fibroblasts. It stimulates the formation of macrophage colonies, enhances antibody-dependent, cell-mediated cytotoxicity by monocytes and macrophages, and inhibits bone resorption by osteoclasts.
Amino Acid Sequence	MGSSHHHHHH SSGLVPRGSH MEEVSEYCSH MIGSGHLQSL QRLIDSQMET SCQITFEFVD QEQLKDPVCY LKKAFLLVQD IMEDTMRFRD NTPNAIAIVQ LQELSLRLKS CFTKDYEEHD KACVRTFYET PLQLLEKVKN VFNETKNLLD KDOWNIFSKNC NNSFAECSSQ DVVTKPDCN
Purity	> 90% by SDS – PAGE.
Form	Liquid. In 20 mM Tris-HCl buffer (pH8.0), 2mM DTT, 10% glycerol.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin Level	< 1.0 EU per 1 µg of protein (determined by LAL method).
Concentration	1 mg/ml (determined by Bradford assay).
Stability	Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.
GENE INFORMATION	
Gene Name	CSF1 colony stimulating factor 1 (macrophage) [Homo sapiens]
Synonyms	CSF1; colony stimulating factor 1 (macrophage); MCSF; MGC31930; colony stimulating factor 1; OTTHUMP00000013363; OTTHUMP00000013364; macrophage colony stimulating factor; M-CSF; CSF-1; Lanimostim; OTTHUMP00000013362
Gene ID	1435
mRNA Refseq	NM_000757
Protein Refseq	NP_000748
MIM	120420
UniProt ID	P09603
Chromosome Location	1p21-p13
Pathway	Cytokine-cytokine receptor interaction; Hematopoietic cell lineage
Function	cytokine activity; growth factor activity; macrophage colony stimulating factor receptor

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



binding; protein binding; protein homodimerization activity

PDB rendering based
on 1hmc.



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA