

## Recombinant Human Cathepsin L2, His-tagged

Cat. No. CTSL2-751H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant human procathepsin V (amino acids 18-334) was fused at the N-terminus to a CD33 signal sequence (amino acids 1-16) and at the C-terminus to a His•Tag sequence and expressed in NSO cells. MW = 38.7 kDa.
<b>Species</b>	Human
<b>Source</b>	Mammalian Cells
<b>Protein Length</b>	18-334 a.a.
<b>Description</b>	<p>Cathepsin L2, also known as CTSL2, Cathepsin V, CATL2, and CTSV, is a member of the peptidase C1 family. It is predominantly expressed in the thymus and testis. Cathepsin V is also expressed in corneal epithelium, and to a lesser extent in conjunctival epithelium and skin. Cathepsin V is a lysosomal cysteine proteinase that may play an important role in corneal physiology. It has about 75% protein sequence identity to murine cathepsin L. The fold of this enzyme is similar to the fold adopted by other members of the papain superfamily of cysteine proteases. Cathepsins K and S are papain-like cysteine proteases with known elastolytic activities, and both enzymes have been identified in macrophages present in plaque areas of diseased blood vessels. Cathepsin V is the third elastolytic cysteine protease which exhibits the most potent elastase activity yet described among human proteases and that Cathepsin V is present in atherosclerotic plaque specimens. Increase of cathepsins V and K activities may accelerate the destruction of the elastin matrix in diseased arteries.</p>
<b>Purity</b>	≥95% by SDS-PAGE.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Form</b>	Frozen liquid. In 800 mM NaCl, 25 mM sodium acetate, 0.8 M NaCl, pH 5.0. Please refer to vial label for lot-specific concentration.
<b>Endotoxins</b>	≤1.0 EU/μg Cathepsin V.
<b>Specific Activity</b>	≥300 units/μg.
<b>Unit Definition</b>	One unit is defined as the amount of enzyme that will cleave 1 pmol the fluorescent substrate Z-LR-AMC per min at 25°C, pH 5.5.
<b>Toxicity</b>	Standard Handling (MSDS available upon request).
<b>Storage</b>	≤-70°C. Avoid freeze/thaw. To activate rhCathepsin V, pre-incubate it at 20 μg/ml in 25 mM NaOAc, 100 mM NaCl, 5 mM DTT, pH 5.5 at room temperature. Following initial thaw aliquot and freeze (-70°C).

## GENE INFORMATION

<b>Gene Name</b>	CTSL2 cathepsin L2 [ Homo sapiens ]
<b>Synonyms</b>	CTSL2; cathepsin L2; CTSU; CTSV; CATL2; MGC125957; EC 3.4.22.43; Cathepsin L2; Cathepsin V; Cathepsin U; UNQ268/PRO305
<b>Gene ID</b>	1515
<b>mRNA Refseq</b>	NM_001333
<b>Protein Refseq</b>	NP_001324
<b>MIM</b>	603308



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>UniProt ID</b>	O60911
<b>Chromosome Location</b>	9q22.2
<b>Pathway</b>	Lysosome
<b>Function</b>	cysteine-type endopeptidase activity; peptidase activity; protein binding

**PDB rendering based on 1fh0.**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA