

Recombinant Human CWC27 293 Cell Lysate

Cat. No. CWC27-7174HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for CWC27 spliceosome-associated protein homolog (<i>S. cerevisiae</i>) (CWC27) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CWC27 CWC27 spliceosome-associated protein homolog (<i>S. cerevisiae</i>) [<i>Homo sapiens</i>]
Official Symbol	CWC27
Synonyms	CWC27; CWC27 spliceosome-associated protein homolog (<i>S. cerevisiae</i>); SDCCAG10, serologically defined colon cancer antigen 10; peptidyl-prolyl cis-trans isomerase CWC27 homolog; NY CO 10; PPIase CWC27; PPIase SDCCAG10; antigen NY-CO-10; peptidyl-prolyl cis-trans isomerase SDCCAG10; serologically defined colon cancer antigen 10; NY-CO-10; SDCCAG10;
Gene ID	10283
mRNA Refseq	NM_005869
Protein Refseq	NP_005860
UniProt ID	Q6UX04
Chromosome Location	5q12.3
Function	isomerase activity; peptidyl-prolyl cis-trans isomerase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA