

## Recombinant Human CX3CL1, Fc-tagged

Cat. No. CX3CL1-209H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	The extracellular domain of human CX3CL1 (aa 25-100) is fused to the N-terminus of the Fc region of human IgG1.
<b>Species</b>	Human
<b>Source</b>	CHO
<b>Description</b>	CX3CL1 [Fractalkine] is expressed on TNF- $\alpha$ and IL-1 activated endothelial cells (ECs) and binds to CX3CR1. CX3CL1 can exist in 2 forms, as membrane-anchored or as a shed 95-kDa glycoprotein. The soluble CX3CL1 has potent chemoattractant activity for T cells and monocytes, and the cell surface bound protein, which is induced on activated primary endothelial cells, promotes strong adhesion of those leukocytes. CX3CL1 may play a role in regulating leukocyte adhesion and migration processes at the endothelium.
<b>Form</b>	Lyophilized from 0.2 $\mu$ m-filtered solution in PBS.
<b>Purity</b>	$\geq$ 98% (SDS-PAGE)
<b>Stability</b>	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
<b>Storage</b>	Short Term Storage: +4°C; Long Term Storage: -20°C. Avoid freeze/thaw cycles.

### GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Gene Name</b>	CX3CL1 chemokine (C-X3-C motif) ligand 1 [ Homo sapiens ]
<b>Official Symbol</b>	CX3CL1
<b>Synonyms</b>	CX3CL1; chemokine (C-X3-C motif) ligand 1; SCYD1, small inducible cytokine subfamily D (Cys X3 Cys), member 1 (fractalkine, neurotactin); fractalkine; ABCD 3; C3Xkine; CXC3; CXC3C; neurotactin; NTN; C-X3-C motif chemokine 1; small-inducible cytokine D1; CX3C membrane-anchored chemokine; small inducible cytokine subfamily D (Cys-X3-Cys), member-1; small inducible cytokine subfamily D (Cys-X3-Cys), member 1 (fractalkine, neurotactin); NTT; SCYD1; ABCD-3;
<b>Gene ID</b>	6376
<b>mRNA Refseq</b>	NM_002996
<b>Protein Refseq</b>	NP_002987
<b>MIM</b>	601880
<b>UniProt ID</b>	P78423
<b>Chromosome Location</b>	16q13
<b>Pathway</b>	Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Direct p53 effectors, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



**Function**

chemokine activity; protein binding; receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA