

Active Recombinant Human CX3CL1 Protein (315 aa)

Cat. No. CX3CL1-283C Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human Fractalkine/CX3CL1 produced in CHO cells is a polypeptide chain containing 315 amino acids. A fully biologically active molecule, rhFractalkine/CX3CL1 has a molecular mass of 50-75 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques.
Species	Human
Source	CHO
ProteinLength	315
Description	<p>Chemokine (C-X3-C motif) ligand 1 (CX3CL1) is a known member of the CX3C chemokine family. It is also commonly known under the names fractalkine (in humans) and neurotactin (in mice). The polypeptide structure of CX3CL1 differs from the typical structure of other chemokines. For example, the spacing of the characteristic N-terminal cysteines is different; there are three amino acids separating the initial pair of cysteines in CX3CL1, while there are none in CC chemokines and only one in CXC chemokines. CX3CL1 is produced as a long protein (with 373-amino acid in humans) with an extended mucin-like stalk and a chemokine domain on top. The mucin-like stalk allows it to bind to the surface of certain cells. Soluble CX3CL1 potently chemoattracts T cells and monocytes, while the cell-bound chemokine promotes strong adhesion of leukocytes to activated endothelial cells, where it is primarily expressed. CX3CL1 can signal through the chemokine receptor CX3CR1.</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	The EC50 value of Human Fractalkine/CX3CL1 on Ca ²⁺ mobilization assay in CHO-K1/Gα15/hCX3CR1 cells (human Gα15 and hCX3CR1 stably expressed in CHO-K1 cells) is less than 1.5 µg/mL.
Molecular Mass	50-75 kDa, observed by reducing SDS-PAGE.
AA Sequence	<p>QHHGVTKCNITCSKMTSKIPVALLIHYQQNQASCGKRAIILETRQHRLFCADPKEQW VKDAMQHLDRQAAALTRNGGTFEKQIGEVKPRTPAAGGMDESVVLEPEATGESS SLEPTPSSQEAQRALGTSPPELPTGVTGSSGTRLPPTPKAQDGGPVGTELFRVPPVS TAATWQNSAPHQPGPSLWAEAKTSEAPSTQDPSTQASTASSPAPEENAPSEGQRV WGQGQSPRPENSLEREEMGPVPAHTDAFQDWGPGSMAHVSVVPVSSEGTPSRE PVASGSWTPKAEPIHATMDPQRLGVLITPVPDAQAATR</p>
Endotoxin	< 0.2 EU/µg, determined by LAL method.
Purity	> 95% as analyzed by SDS-PAGE.
Storage	Lyophilized recombinant Human Fractalkine/CX3CL1 remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, Human Fractalkine/CX3CL1 should be stable up to 1 week at 4 centigrade or up to 3 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 µg/mL.

GENE INFORMATION

Gene Name CX3CL1 chemokine (C-X3-C motif) ligand 1 [Homo sapiens]

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	CX3CL1
Synonyms	CX3CL1; chemokine (C-X3-C motif) ligand 1; SCYD1, small inducible cytokine subfamily D (Cys X3 Cys), member 1 (fractalkine, neurotactin); fractalkine; ABCD 3; C3Xkine; CXC3; CXC3C; neurotactin; NTN; C-X3-C motif chemokine 1; small-inducible cytokine D1; CX3C membrane-anchored chemokine; small inducible cytokine subfamily D (Cys-X3-Cys), member-1; small inducible cytokine subfamily D (Cys-X3-Cys), member 1 (fractalkine, neurotactin); NTT; SCYD1; ABCD-3;
Gene ID	6376
mRNA Refseq	NM_002996
Protein Refseq	NP_002987
MIM	601880
UniProt ID	P78423

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA