

Active Recombinant Human Chemokine (C-X-C Motif) Ligand 1, MlgG2a Fc-tagged

Cat. No. CXCL1-222H Lot. No. (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human CXCL1 (AAH11976.1)(Ala35-Asn107) is fused to the N-terminus of the Fc region of mouse IgG2a was expressed in CHO cell.
Species	Human
Source	CHO
ProteinLength	35-107 a.a.
Description	This gene encodes a member of the CXC subfamily of chemokines. The encoded protein is a secreted growth factor that signal through the G-protein coupled receptor, CXC receptor 2. This protein plays a role in inflammation and as a chemoattractant for neutrophils. Aberrant expression this protein is associated with the growth and progression of certain tumors. A naturally occurring processed form of this protein has increased chemotactic activity. Alternate splicing results in coding and non-coding variants of this gene. A pseudogene of this gene is found on chromosome 4
Form	Lyophilized from 0.2µm-filtered solution in PBS.
Bio-activity	Measured by its ability to induce myeloperoxidase.
Molecular Mass	34KDa (monomer)
AA Sequence	Ala35-Asn107

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	<0.06 eu/μg="" as="" determined="" by="" la="">
Purity	>98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100μg/ml in sterile PBS.
Warning	Avoid freeze/thaw cycles.
GENE INFORMATION	
Gene Name	CXCL1 chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, alpha) [Homo sapiens]
Official Symbol	CXCL1
Synonyms	CXCL1; chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, alpha); fibroblast secretory protein , FSP, GRO1, GRO1 oncogene (melanoma growth stimulating activity, alpha) , MGSA; growth-regulated alpha protein; GROa; MGSA a; NAP 3; SCYB1; MGSA alpha; GRO-alpha(1-73); C-X-C motif chemokine 1; fibroblast secretory protein; neutrophil-activating protein 3; melanoma growth stimulatory activity alpha; GRO1 oncogene (melanoma growth-stimulating activity); GRO1 oncogene (melanoma growth stimulating activity, alpha); FSP; GRO1; MGSA; NAP-3; MGSA-a;
Gene ID	2919
mRNA Refseq	NM_001511

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_001502
MIM	155730
UniProt ID	P09341
Chromosome Location	4q13.3
Pathway	Amoebiasis, organism-specific biosystem; Amoebiasis, conserved biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem;
Function	chemokine activity; enzyme activator activity; growth factor activity; receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA