

Recombinant Equine Chemokine (C-X-C motif) Ligand 10

Cat. No. CXCL10-142E Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Equine CXCL10 protein was expressed in Yeast and purified by using Ion-exchange chromatography, 9.4 kDa (83 aa).
Species	Equine
Source	Yeast
Description	Chemokine (C-X-C motif) ligand 10 (CXCL10) or IP-10 is a small cytokine belonging to the CXC chemokine family that is also known as 10 kDa interferon-gamma-induced protein (γ -IP10 or IP-10). CXCL10 is secreted by several cell types in response to IFN- γ . These cell types include monocytes, endothelial cells and fibroblasts.
Sequences Of Amino Acids	IPLSRTARCT CINISDRPIP PRSLEKLEMI PASQSCQRVE IIATMKKNGE KRCLNPESKT VKNLLKAISK QRSKRSPRTL REV.
Formulation/Reconstitution	Lyophilized. Reconstitute with sterile phosphate-buffered saline containing at least 0.1% carrier protein.
Purity	>95% as visualized by SDS-PAGE analysis.
Purification	Ion-exchange chromatography.
Biological Activity	In testing.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Stable for up to twelve months from date of receipt at -20°C. Stable for at least 3 months when stored in working aliquots with a carrier protein at -20°C. Avoid repeated freeze/thaw cycles.
Pathways	Chemokine signaling pathway; Cytokine-cytokine receptor interaction; Cytosolic DNA-sensing pathway; RIG-I-like receptor signaling pathway; Toll-like receptor signaling pathway

GENE INFORMATION

Gene Name	CXCL10 chemokine (C-X-C motif) ligand 10 [<i>Equus caballus</i>]
Synonyms	CXCL10; chemokine (C-X-C motif) ligand 10
Gene ID	100050993
mRNA Refseq	NM_001114940
Protein Refseq	NP_001108412
Chromosome Location	3

**PDB rendering based on
1o7z**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA