

## Recombinant Mouse CXCL10 (IP-10)

Cat. No. CXCL10-71M Lot. No. (See product label)

### SPECIFICATION

**Product Overview** Mouse IP-10/CXCL10 was produced in yeast and therefore does not have endotoxin, is naturally folded, and post-translationally modified.

**Species** Mouse

**Source** Yeast

**Description** C-X-C motif chemokine 10 (CXCL10) also known as Interferon gamma-induced protein 10 (IP-10) or small-inducible cytokine B10 is a member of the C-X-C chemokine family. CXCL10 (IP-10) is secreted by several cell types in response to IFN-gamma. These cell types include monocytes, endothelial cells and fibroblasts. CXCL10 (IP-10) has been attributed to several roles, such as chemoattraction for monocytes/macrophages, T cells, NK cells, and dendritic cells, promotion of T cell adhesion to endothelial cells, antitumor activity, and inhibition of bone marrow colony formation and angiogenesis. There have been 17 different C-X-C chemokines described in mammals, that are subdivided into two categories, those with a specific amino acid sequence (or motif) of glutamic acid-leucine-arginine (or ELR for short) immediately before the first cysteine of the C-X-C motif (ELR-positive), and those without an ELR motif (ELR-negative). ELR-positive C-X-C chemokines specifically induce the migration of neutrophils, and interact with chemokine receptors CXCR1 and CXCR2. C-X-C chemokines that lack the ELR motif are chemoattractant for lymphocytes. CXCL10 (IP-10) elicits its effects by binding to the cell surface chemokine receptor CXCR3.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Form</b>	Lyophilized
<b>Molecular Mass</b>	8.7 kDa
<b>AA Sequence</b>	IPLARTVRCN CIHIDDGPVR MRAIGKLEII PASLSCPRVE IIATMKKNDE QRCLNPESKT IKNLMKAFSQ KRSKRAP (77)
<b>Applications</b>	The mouse IP-10/CXCL10 protein can be used in cell culture, as an IP-10/CXCL10 ELISA Standard, and as a Western Blot Control.
<b>Storage</b>	-20 C
<b>GENE INFORMATION</b>	
<b>Gene Name</b>	<a href="#">Cxcl10 chemokine (C-X-C motif) ligand 10 [ Mus musculus ]</a>
<b>Official Symbol</b>	CXCL10
<b>Synonyms</b>	CXCL10; chemokine (C-X-C motif) ligand 10; C-X-C motif chemokine 10; gamma-IP10; interferon activated gene 10; small-inducible cytokine B10; interferon-gamma induced protein CRG-2; interferon-gamma-induced protein CRG-2; 10 kDa interferon gamma-induced protein; small inducible cytokine B subfamily (Cys-X-Cys), member 10; C7; IP10; CRG-2; INP10; IP-10; Ifi10; mob-1; Scyb10; gIP-10;
<b>Gene ID</b>	<a href="#">15945</a>
<b>mRNA Refseq</b>	<a href="#">NM_021274</a>
<b>Protein Refseq</b>	<a href="#">NP_067249</a>
<b>Pathway</b>	Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Cytosolic DNA-sensing pathway, organism-specific biosystem;

**Function**

chemokine activity; cytokine activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA