

Active Recombinant Human Chemokine (C-X-C Motif) Ligand 11, HIgG1 Fc-tagged, mutant

Cat. No. CXCL11-166H Lot. No. (See product label)

SPECIFICATION

Product Overview

The extracellular domain of human CXCL11 (AAH05292.1)(Phe22-Phe94) is fused to the N-terminus of the Fc region of a mutant human IgG1 was expressed in CHO cell.

Species

Human

Source

CHO

ProteinLength

22-94 a.a.

Description

Chemokines are a group of small (approximately 8 to 14 kD), mostly basic, structurally related molecules that regulate cell trafficking of various types of leukocytes through interactions with a subset of 7-transmembrane, G protein-coupled receptors. Chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis. Chemokines are divided into 2 major subfamilies, CXC and CC. This gene is a CXC member of the chemokine superfamily. Its encoded protein induces a chemotactic response in activated T-cells and is the dominant ligand for CXC receptor-3. The gene encoding this protein contains 4 exons and at least three polyadenylation signals which might reflect cell-specific regulation of expression. IFN-gamma is a potent inducer of transcription of this gene.

Form

Lyophilized from 0.2µm-filtered solution in PBS.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Bio-activity	Measured by its ability to chemoattract human peripheral blood lymphocytes (PBL).
Molecular Mass	34KDa (monomer)
AA Sequence	Phe22-Phe94
Endotoxin	<0.06 eu/μg="" as="" determined="" by="" lal="" test="">
Purity	>98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100μg/ml in sterile PBS.
Warning	Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name	CXCL11 chemokine (C-X-C motif) ligand 11 [Homo sapiens]
Official Symbol	CXCL11
Synonyms	CXCL11; chemokine (C-X-C motif) ligand 11; SCYB9B, SCYB11, small inducible cytokine subfamily B (Cys X Cys), member 11; C-X-C motif chemokine 11; b R1; H174; I TAC; IP 9; beta-R1; small inducible cytokine B11; small-inducible cytokine B11; interferon gamma-inducible protein 9; interferon-inducible T-cell alpha chemoattractant; small inducible cytokine subfamily B (Cys-X-Cys), member 11; small inducible cytokine subfamily B (Cys-X-Cys), member 9B; IP9; IP-9; b-R1; I-TAC; SCYB11; SCYB9B; MGC102770;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	6373
mRNA Refseq	NM_005409
Protein Refseq	NP_005400
MIM	604852
UniProt ID	O14625
Chromosome Location	4q21
Pathway	CXCR3-mediated signaling events, organism-specific biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;
Function	chemokine activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA