

Active Recombinant Human CXCL11 Protein (73 aa)

Cat. No. CXCL11-174C Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human I-TAC/CXCL11 produced in HEK293 cells is a single non-glycosylated polypeptide chain containing 73 amino acids. A fully biologically active molecule, rHI-TAC/CXCL11 has a molecular mass of 8.3 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques.
Species	Human
Source	HEK293
ProteinLength	73
Description	<p>Chemokine (C-X-C motif) ligand 11(CXCL11), also known as I-TAC and B-R1, is a small cytokine belonging to the CXC chemokine family that is also called Interferon-inducible T-cell alpha chemoattractant (I-TAC) and Interferon-gamma-inducible protein 9 (IP-9). This chemokine elicits its effects on target cells by interacting with chemokine receptor CXCR3 having a higher affinity than other ligands for this receptor such as CXCL9 and CXCL10. CXCL11 is chemotactic for activated T cells. The gene encoding CXCL11 has been mapped to chromosome 4. CXCL11 cDNA encodes a 94 amino acid residue precursor protein with a 21 amino acid residue putative signal sequence, which is cleaved to form the mature 73 amino acid residue protein. CXCL11 shares 36% and 37% amino acid sequence homology with IP-10 and MIG (two other known human non-ELR CXC chemokines), respectively. Mouse CXCL11 exhibits 68% sequence homology with human CXCL11.</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	The EC50 value of human I-TAC/CXCL11 on Ca ²⁺ mobilization assay in CHO-K1/Ga15/hCXCR3 cells (human Ga15 and human CXCR3 stably expressed in CHO-K1 cells) is less than 0.5 µg/mL.
Molecular Mass	8.3 kDa, observed by reducing SDS-PAGE.
AA Sequence	FPMFKRGRCLCIGPGVKAVKVADIEKASIMYPSNNCDKIEVIITLKENKGQRCLNPKS KQARLIIKKVERKNF
Endotoxin	< 0.2 EU/µg, determined by LAL method.
Purity	> 98% as analyzed by SDS-PAGE.
Storage	Lyophilized recombinant human I-TAC/CXCL11 remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, human CXCL11/I-TAC should be stable up to 1 week at 4 centigrade or up to 2 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 µg/mL.

GENE INFORMATION

Gene Name	CXCL11 chemokine (C-X-C motif) ligand 11 [Homo sapiens]
Official Symbol	CXCL11
Synonyms	CXCL11; chemokine (C-X-C motif) ligand 11; SCYB9B, SCYB11, small inducible cytokine subfamily B (Cys X Cys), member 11; C-X-C motif chemokine 11; b R1;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



H174; I TAC; IP 9; beta-R1; small inducible cytokine B11; small-inducible cytokine B11; interferon gamma-inducible protein 9; interferon-inducible T-cell alpha chemoattractant; small inducible cytokine subfamily B (Cys-X-Cys), member 11; small inducible cytokine subfamily B (Cys-X-Cys), member 9B; IP9; IP-9; b-R1; I-TAC; SCYB11; SCYB9B; MGC102770;

Gene ID [6373](#)

mRNA Refseq [NM_005409](#)

Protein Refseq [NP_005400](#)

MIM [604852](#)

UniProt ID [O14625](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA