

Active Recombinant Human CXCL6, MlgG2a Fc-tagged

Cat. No. CXCL6-123H Lot. No. (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human CXCL6 (AAH13744.1)(Val40-Asn114) is fused to the N-terminus of the Fc region of mouse IgG2a was expressed in CHO cell.
Species	Human
Source	CHO
ProteinLength	40-114 a.a.
Form	Lyophilized from 0.2µm-filtered solution in PBS.
Bio-activity	Measured by its ability to induce myeloperoxidase.
Molecular Mass	34KDa (monomer)
AA Sequence	Val40-Asn114
Endotoxin	<0.06 eu/µg="" as="" determined="" by="" lal="" test.="">
Purity	>98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100µg/ml in sterile PBS.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Warning Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name CXCL6 chemokine (C-X-C motif) ligand 6 (granulocyte chemotactic protein 2) [Homo sapiens]

Official Symbol CXCL6

Synonyms CXCL6; chemokine (C-X-C motif) ligand 6 (granulocyte chemotactic protein 2); SCYB6, small inducible cytokine subfamily B (Cys X Cys), member 6 (granulocyte chemotactic protein 2); C-X-C motif chemokine 6; CKA 3; GCP 2; chemokine alpha 3; small-inducible cytokine B6; granulocyte chemotactic protein 2; Small inducible cytokine subfamily B (Cys-X-Cys), member b; small inducible cytokine subfamily B (Cys-X-Cys), member 6 (granulocyte chemotactic protein 2); GCP2; CKA-3; GCP-2; SCYB6;

Gene ID 6372

mRNA Refseq NM_002993

Protein Refseq NP_002984

MIM 138965

UniProt ID P80162

Chromosome Location 4q13.3

Pathway Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; G alpha (i) signalling events, organism-specific biosystem;

Function

chemokine activity; heparin binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA