

Active Recombinant Human CXCL8 Protein (77 aa)

Cat. No. CXCL8-350C Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human Interleukin-8 (IL-8, 77aa)/CXCL8 produced in E. coli is a single non-glycosylated polypeptide chain containing 77 amino acids. A fully biologically active molecule, rhIL-8(77aa)/CXCL8 has a molecular mass of 8.9 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques.
Species	Human
Source	E.coli
ProteinLength	77
Description	Interleukin-8 is one of the first discovered chemokines and belongs to the CXCL family, in which the first two conserved cysteines are separated by one residue. In vivo, IL-8 exists in two forms: a 77 a.a. protein produced by endothelial cells, and the more active 72 a.a. protein produced by monocytes. The receptors for IL-8 are the seven-helical G-protein coupled receptors CXCR1 and CXCR2, exclusively expressed on neutrophils. The functions of IL-8 are to induce rapid changes in cell morphology, activate integrins, and release the granule contents of neutrophils. Thus, IL-8 can enhance the antimicrobial actions of defense cells.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	The EC ₅₀ value of human IL-8(77aa) on Ca ²⁺ mobilization assay in CHO-K1/Gα15/hCXCR1 cells (human Gα15 and human CXCR1 stably expressed in CHO-K1 cells) is less than 150 ng/mL.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	8.9 kDa, observed by reducing SDS-PAGE.
AA Sequence	AVLPRSAKELRCQCIKTYSKPFHPKFIKELRVIESGPHCANTEIIVKLSDGRELCLDPK ENWVQRVVEKFLKRAENS
Endotoxin	< 0.2 EU/μg, determined by LAL method.
Purity	> 95% as analyzed by SDS-PAGE.
Storage	Lyophilized recombinant human Interleukin-8 (IL-8)(77aa)/CXCL8 remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, recombinant human Interleukin-8 (IL-8)(77aa)/CXCL8 should be stable up to 1 week at 4 centigrade or up to 2 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 μg/mL.

GENE INFORMATION

Gene Name	CXCL8 C-X-C motif chemokine ligand 8 [Homo sapiens (human)]
Official Symbol	CXCL8
Synonyms	CXCL8; C-X-C motif chemokine ligand 8; IL8; NAF; GCP1; LECT; LUCT; NAP1; GCP-1; LY NAP; MDNCF; MONAP; NAP-1; SCYB8; interleukin-8; T-cell chemotactic factor; alveolar macrophage chemotactic factor I; beta endothelial cell-derived neutrophil activating peptide; beta-thromboglobulin-like protein; chemokine (C-X-C motif) ligand 8; emoctakin; granulocyte chemotactic protein 1; interleukin 8; lung giant cell carcinoma-derived chemotactic protein; lymphocyte derived neutrophil activating peptide; monocyte-derived neutrophil chemotactic factor; monocyte-derived

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



neutrophil-activating peptide; neutrophil-activating peptide 1; small inducible cytokine subfamily B, member 8; tumor necrosis factor-induced gene 1

Gene ID 3576

mRNA Refseq NM_000584

Protein Refseq NP_000575

MIM 146930

UniProt ID P10145

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA