

## Recombinant Mouse CXCL9 (MIG)

**Cat. No.** CXCL9-73M    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Mouse CXCL9/MIG was produced in yeast and therefore does not have endotoxin, is naturally folded, and post-translationally modified.
<b>Species</b>	Mouse
<b>Source</b>	Yeast
<b>Description</b>	Chemokine (C-X-C motif) ligand 9 (CXCL9) is a small cytokine belonging to the CXC chemokine family. It is also commonly known as Monokine Induced by Gamma interferon (MIG). CXCL9 (MIG) is a T-cell chemoattractant, which is induced by IFN-gamma. It is closely related to two other CXC chemokines -CXCL10 and CXCL11. There have been 17 different C-X-C chemokines described in mammals, that are subdivided into two categories, those with a specific amino acid sequence (or motif) of glutamic acid-leucine-arginine (or ELR for short) immediately before the first cysteine of the C-X-C motif (ELR-positive), and those without an ELR motif (ELR-negative). ELR-positive C-X-C chemokines specifically induce the migration of neutrophils, and interact with chemokine receptors CXCR1 and CXCR2. C-X-C chemokines that lack the ELR motif are chemoattractant for lymphocytes. CXCL9 (MIG) elicits its effects by binding to the cell surface chemokine receptor CXCR3.
<b>Form</b>	Lyophilized
<b>Molecular Mass</b>	12.2 kDa
<b>AA Sequence</b>	TLVIRNARCS CISTSRGTIH YKSLKDLKQF APSPNCNKTE IIATLKNGDQ

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

TCLDPDSANV KKLMKEWEKK INQKKKQKRG KKHQKNMKNR KPKTPQSRRR  
SRKTT (105)

**Applications**

The mouse CXCL9/MIG protein can be used in cell culture, as a CXCL9/MIG ELISA Standard, and as a Western Blot Control.

**Storage**

-20 C

## GENE INFORMATION

**Gene Name**

Cxcl9 chemokine (C-X-C motif) ligand 9 [ *Mus musculus* ]

**Official Symbol**

CXCL9

**Synonyms**

CXCL9; chemokine (C-X-C motif) ligand 9; C-X-C motif chemokine 9; M119; protein m119; small-inducible cytokine B9; gamma interferon-induced monokine; gamma-interferon-induced monokine; monokine induced by gamma interferon; monokine induced by interferon-gamma; small inducible cytokine B subfamily (Cys-X-Cys), member 9; CMK; Mig; MuMIG; Scyb9; crg-10; BB139920;

**Gene ID**

17329

**mRNA Refseq**

NM\_008599

**Protein Refseq**

NP\_032625

**Pathway**

Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; G alpha (i) signalling

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



events, organism-specific biosystem;

**Function**

cytokine activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA