

Recombinant Human CXCR4 293 Cell Lysate

Cat. No. CXCR4-7161HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for chemokine (C-X-C motif) receptor 4 (CXCR4), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name CXCR4 chemokine (C-X-C motif) receptor 4 [Homo sapiens]

Official Symbol CXCR4

Synonyms

CXCR4; chemokine (C-X-C motif) receptor 4; chemokine (C X C motif), receptor 4 (fusin); C-X-C chemokine receptor type 4; CD184; D2S201E; fusin; HM89; HSY3RR; LESTR; NPY3R; NPYR; NPYY3R; CXC-R4; CXCR-4; CD184 antigen; SDF-1 receptor; neuropeptide Y receptor Y3; seven transmembrane helix receptor; stromal cell-derived factor 1 receptor; lipopolysaccharide-associated protein 3; seven-transmembrane-segment receptor, spleen; leukocyte-derived seven transmembrane domain receptor; leukocyte-derived seven-transmembrane-domain receptor; FB22; LAP3; LCR1; WHIM; NPYRL;

Gene ID 7852

mRNA Refseq NM_001008540

Protein Refseq NP_001008540

MIM 162643

UniProt ID P61073

Chromosome Location 2q21

Pathway Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Binding and entry of HIV virion, organism-specific biosystem; CXCR4-mediated signaling events, organism-specific biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem;

Function

C-X-C chemokine receptor activity; G-protein coupled receptor activity; actin binding; coreceptor activity; myosin light chain binding; protein binding; receptor activity; signal transducer activity; ubiquitin binding; ubiquitin protein ligase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA