

Recombinant Human CYP27A1 293 Cell Lysate

Cat. No. CYP27A1-7119HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for cytochrome P450, family 27, subfamily A, polypeptide 1 (CYP27A1), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [CYP27A1 cytochrome P450, family 27, subfamily A, polypeptide 1 \[Homo sapiens \]](#)

Official Symbol CYP27A1

Synonyms

CYP27A1; cytochrome P450, family 27, subfamily A, polypeptide 1; CYP27, cytochrome P450, subfamily XXVIIA (steroid 27 hydroxylase, cerebrotendinous xanthomatosis), polypeptide 1; sterol 26-hydroxylase, mitochondrial; cerebrotendinous xanthomatosis; CP27; CTX; cytochrome P450 27; sterol 27-hydroxylase; cytochrome P-450C27/25; vitamin D(3) 25-hydroxylase; cholestanetriol 26-monooxygenase; 5-beta-cholestane-3-alpha,7-alpha,12-alpha-triol 27-hydroxylase; 5-beta-cholestane-3-alpha, 7-alpha, 12-alpha-triol 26-hydroxylase; 5-beta-cholestane-3-alpha, 7-alpha, 12-alpha-triol 27-hydroxylase; cytochrome P450, subfamily XXVIIA (steroid 27-hydroxylase, cerebrotendinous xanthomatosis), polypeptide 1; CYP27;

Gene ID [1593](#)

mRNA Refseq [NM_000784](#)

Protein Refseq [NP_000775](#)

MIM [606530](#)

UniProt ID [Q02318](#)


Chromosome 2q33-qter

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	Bile acid and bile salt metabolism, organism-specific biosystem; Bile acid biosynthesis, cholesterol =>cholate, organism-specific biosystem; Bile acid biosynthesis, cholesterol => cholate, conserved biosystem; Biological oxidations, organism-specific biosystem; Cytochrome P450 - arranged by substrate type, organism-specific biosystem;
Function	cholestanetriol 26-monooxygenase activity; cholesterol 26-hydroxylase activity; electron carrier activity; heme binding; metal ion binding; oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen; steroid hydroxylase activity; vitamin D3 25-hydroxylase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA