

Recombinant Human CYP2C8 protein, His&Myc-tagged

Cat. No. CYP2C8-2794H Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant Human CYP2C8 protein(P10632)(27-490aa), fused to N-terminal His tag and C-terminal Myc tag, was expressed in E. coli.

Species Human

Source E.coli

ProteinLength 27-490aa

Form If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol.
If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Molecular Mass 60.1 kDa

AA Sequence

```
RKLPPGPTPLPIIGNMLQIDVKDICKSFTNFSKVYGPVFTVYFGMNPVVFHGYEAVK
EALIDNGEEFSGRGNPISQRITKGLGISSNGKRWKEIRRFSLTTLRNFGMGKRSIED
RVQEEAHCLVEELRKTASPCDPTFILGCAPCNVICSVVFQKRFDYKQDNFLTLMKR
FNENFRILNSPWIQVCNPFLLIDCFPGTHNKVLKNVALTRSYIREKVKEHQASLDVN
NPRDFIDCFLIKMEQEKDNQKSEFNENLVGTVADLFVAGTETTSTTLRYGLLLLLLKH
EVTAKVQEEIDHVIGRHRSPCMQDRSHMPYTDVAVVHEIQRYSDLVPTGVPHAVTTD
TKFRNYLIPKGTTIMALLTSVLHDDKEFPNPNIFDPGHFLDKNGNFKKSDFMPFSAG
KRICAGEGLARMELFLFLTTILQNFNLKSVDDLKLNLTAVTKGIVSLPPSYQICFIPV
```

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	Greater than 85% as determined by SDS-PAGE.
Storage	Store at -20°C/-80°C upon receipt, aliquoting is necessary for mutiple use. Avoid repeated freeze-thaw cycles.
Reconstitution	Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%.

GENE INFORMATION

Gene Name	CYP2C8 cytochrome P450, family 2, subfamily C, polypeptide 8 [Homo sapiens]
Official Symbol	CYP2C8
Synonyms	CYP2C8; cytochrome P450, family 2, subfamily C, polypeptide 8; cytochrome P450, subfamily IIC (mephenytoin 4 hydroxylase), polypeptide 8; cytochrome P450 2C8; CPC8; P450 form 1; cytochrome P450 IIC2; cytochrome P450 MP-12; cytochrome P450 MP-20; cytochrome P450 form 1; microsomal monooxygenase; xenobiotic monooxygenase; s-mephenytoin 4-hydroxylase; flavoprotein-linked monooxygenase; cytochrome P450, subfamily IIC (mephenytoin 4-hydroxylase), polypeptide 8; CYP1IC8; MP-12/MP-20;
Gene ID	1558
mRNA Refseq	NM_000770
Protein Refseq	NP_000761
UniProt ID	P10632

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA