

Active Recombinant Human CYP2D6 Protein

Cat. No. CYP2D6-78H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human CYP2D6 protein and CYP-reductase without tag were coexpressed in E. coli.
Species	Human
Source	E.coli
Description	<p>This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is known to metabolize as many as 25% of commonly prescribed drugs. Its substrates include antidepressants, antipsychotics, analgesics and antitussives, beta adrenergic blocking agents, antiarrhythmics and antiemetics. The gene is highly polymorphic in the human population; certain alleles result in the poor metabolizer phenotype, characterized by a decreased ability to metabolize the enzyme's substrates. Some individuals with the poor metabolizer phenotype have no functional protein since they carry 2 null alleles whereas in other individuals the gene is absent. This gene can vary in copy number and individuals with the ultrarapid metabolizer phenotype can have 3 or more active copies of the gene. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.</p>
Bio-activity	<p>Cytochrome c Reductase Activity: 310 nmol/min/mg protein P450 Activity Data(Unless otherwise stated, all assays are carried out at 37</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

centigrade in 50 mM potassium phosphate buffer (pH 7.4) containing 5mM magnesium chloride and an NADPH generating system [5 mM glucose 6-phosphate, 1 mM NADP+, 1 U/ml glucose 6-phosphate dehydrogenase]

Activity: Dextromethorphan O-demethylase

Linear with time up to: 30 min(Determined at 0.4 μ M substrate, using a P450 concentration of 5 pmol/ml)

Linear with P450 up to: 30 pmol/ml(Determined at 0.4 μ M substrate, using an incubation time of 5 min)

Vmax: 2.7 min⁻¹

Km: 0.85 μ M

Usage For laboratory (research) purposes only

Storage Store at -80 centigrade. Avoid frequent temperature changes. Thaw on ice.

Concentration

P450 Concentration: 5.7 nmol/ml
 Protein Concentration: 8.7 mg/ml
 Specific P450 Content: 655 pmol/mg protein

Storage Buffer 50 mM Tris-acetate (pH 7.6), 250 mM sucrose, 0.25 mM EDTA

GENE INFORMATION

Gene Name CYP2D6 cytochrome P450 family 2 subfamily D member 6 [Homo sapiens (human)]

Official Symbol CYP2D6

Synonyms CYP2D6; cytochrome P450 family 2 subfamily D member 6; CPD6; CYP2D; CYP2DL1; CYPIID6; P450C2D; P450DB1; CYP2D7AP; CYP2D7BP; CYP2D7P2; CYP2D8P2; P450-DB1; cytochrome P450 2D6; cholesterol 25-hydroxylase; cytochrome P450, family 2, subfamily D, polypeptide 6; cytochrome P450, family 2,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

subfamily D, polypeptide 7 pseudogene 2; cytochrome P450, family 2, subfamily D, polypeptide 8 pseudogene 2; cytochrome P450, subfamily II (debrisoquine, sparteine, etc., -metabolising), polypeptide 7 pseudogene 2; cytochrome P450, subfamily IID (debrisoquine, sparteine, etc., -metabolising), polypeptide 8 pseudogene 2; cytochrome P450, subfamily IID (debrisoquine, sparteine, etc., -metabolizing), polypeptide 6; cytochrome P450, subfamily IID (debrisoquine, sparteine, etc., -metabolizing)-like 1; cytochrome P450-DB1; debrisoquine 4-hydroxylase; flavoprotein-linked monooxygenase; microsomal monooxygenase; nonfunctional cytochrome P450 2D6; xenobiotic monooxygenase

Gene ID [1565](#)

mRNA Refseq [NM_000106](#)

Protein Refseq [NP_000097](#)

MIM [124030](#)

UniProt ID [P10635](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA