

Recombinant Murine Chemokine (C-C Motif) Ligand 4

Cat. No. Ccl4-105M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant murine MIP-1 beta produced in <i>E.coli</i> is a 7.8 kDa protein containing 69 amino acid residues, including the four highly conserved cysteine residues present in CC chemokines.
Species	Mouse
Source	E.coli
Description	Both MIP-1 alpha and MIP-1 beta are structurally and functionally related CC chemokines. They participate in the host response to invading bacterial, viral, parasite and fungal pathogens by regulating the trafficking and activation state of selected subgroups of inflammatory cells e.g. macrophages, lymphocytes and NK cells. While both MIP-1 alpha and MIP-1 beta exert similar effects on monocytes their effect on lymphocytes differ; with MIP-1 alpha selectively attracting CD8+ lymphocytes and MIP-1 beta selectively attracting CD4+ lymphocytes. Additionally, MIP-1 alpha and MIP-1 beta have also been shown to be potent chemoattractants for B cells, eosinophils and dendritic cells. Both human and murine MIP-1 alpha and MIP-1 beta are active on human and murine hematopoietic cells.
Purity	> 95% by SDS-PAGE.
Endotoxin level	< 0.1 ng per µg of MIP-1 beta.
Stabilizer	None.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Formulation	Lyophilized.
Biologically Activity	Determined by its ability to chemoattract human monocytes using a concentration range of 20.0-100.0 ng/ml.
Reconstitution	The murine MIP-1 beta should be reconstituted in water to a concentration of 0.1-1.0 mg/ml. This solution can be diluted in water or other buffer solutions or stored at -20°C.
Stability	The lyophilized protein is stable for a few weeks at room temperature, but best stored at -20°C. Reconstituted murine MIP-1 beta should be stored in working aliquots at -20°C. Avoid repeated freeze-thaw cycles.

GENE INFORMATION

Gene Name	Ccl4 chemokine (C-C motif) ligand 4 [Mus musculus]
Synonyms	Ccl4; chemokine (C-C motif) ligand 4; Act-2; Mip1b; Scya4; MIP-1B; AT744.1
Gene ID	20303
mRNA Refseq	NM_013652
Protein Refseq	NP_038680
UniProt ID	P14097
Chromosome Location	47.6 cM
pathway	Chemokine signaling pathway; Cytokine-cytokine receptor interaction; Toll-like

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

receptor signaling pathway

Function

chemokine activity; cytokine activity; protein binding

PDB rendering based
on 1hum.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA