

Active Recombinant Mouse Chemokine (C-C motif) Ligand 9

Cat. No. Ccl9-400M Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant mouse Ccl9 protein was expressed in E. coli. Gly50-Gln122.
Species	Mouse
Source	E.coli
Protein Length	50-122 a.a.
Description	Mouse Ccl9 (also MIP1 γ and MRP2) is an 11 kDa, secreted, monomeric polypeptide that belongs to the β (or CC) intercrine family of chemokines. Based on its activity and amino acid (aa) sequence, it is further classified as a member of the NC6 or six cysteine containing CC subfamily of chemokines.
Form	Lyophilized from a 0.2 μ m filtered solution in PBS with BSA as a carrier protein.
N-terminal Sequence	Gly50
Molecular Weight	8.4 kDa
Activity	Measured by its ability to chemoattract human CCR1 transfected BaF3 mouse pro-B cells. The ED50 for this effect is typically 0.2-1 ng/mL.
Endotoxin Level	<1.0 EU per 1 μ g of the protein by the LAL method.
Purity	>97%, by SDS-PAGE under reducing conditions and visualized by silver stain.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Reconstitution	Reconstitute at 100µg/mL in sterile PBS that containing at least 0.1% human or bovine serumalbumin.
Storage	Use amannual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 1month, 2 to 8°C under sterile conditions after reconstitution. 3 months, -20 to-70°C under sterile conditions after reconstitution.
GENE INFORMATION	
Gene Name	Ccl9 chemokine (C-C motif)ligand 9 [Mus musculus]
Official Symbol	Ccl9
Synonyms	Ccl9; chemokine(C-C motif) ligand 9; CCF18; MRP-2; Scya9; Scya10; OTTMUSP00000000947;MIP-1 gamma; C-C motif chemokine 9; small induciblecytokine A9; small inducible cytokine A10; macrophage inflammatory protein1-gamma; macrophage inflammatory protein-ratated protein 2; MIP-1-gamma;Mrp2; Small-inducible cytokine A9
Gene ID	20308
mRNA Refseq	NM_011338
Protein Refseq	NP_035468
UniProt ID	P51670
Chromosome Location	11
Pathway	Chemokine signalingpathway; Class A/1 (Rhodopsin-like receptors); Cytokine-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



cytokine receptorinteraction; GPCR ligand binding; GPCR downstream signaling; G alpha (i)signaling events; G alpha (q) signaling events; Peptide ligand-bindingreceptors; Signal Transduction; Signaling by GPCR

Function

chemokineactivity; cytokine activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA