

Active Recombinant Mouse Cd27 Protein, His & Fc-tagged

Cat. No. Cd27-840M Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant extracellular domain (Met 1-Arg 182) of mouse CD27 (NP_001028298.1) precursor was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Species	Mouse
Source	HEK293
ProteinLength	1-182 a.a.
Predicted N Terminal	Thr 21
Form	Lyophilized from sterile PBS, pH 7.4, 5%~8% trehalose and mannitol.
Bio-activity	Measured by its binding ability in a functional ELISA. Immobilized recombinant mouse CD27 at 2 µg/ml (100 µl/well) can bind biotinylated human CD70 with a linear range of 0.39-12.5 ng/ml.
Molecular Mass	The recombinant mouse CD27/Fc is a disulfide-linked homodimer after removal of the signal peptide. The reduced monomer consists of 410 amino acids and has a predicted molecular mass of 46.2 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rm CD27/Fc monomer is approximately 60-65 kDa due to glycosylation.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	>94 % as determined by SDS-PAGE.
Stability	Samples are stable for up to twelve months from date of receipt at -70°C.
Storage	Store it under sterile conditions at -20°C~-70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.25 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	Cd27 CD27 antigen [Mus musculus]
Official Symbol	Cd27
Synonyms	CD27; CD27 antigen; CD antigen 27; CD27L receptor; T-cell activation antigen CD27; tumor necrosis factor receptor superfamily member 7; tumor necrosis factor receptor superfamily, member 7; S152; Tp55; Tnfrsf7;
Gene ID	21940
mRNA Refseq	NM_001033126
Protein Refseq	NP_001028298
MIM	
UniProt ID	
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

binding; cysteine-type endopeptidase inhibitor activity involved in apoptotic process;
protein binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA