

Recombinant Mouse Cebpb Protein, MYC/DDK-tagged

Cat. No. Cebpb-876M Lot. No. (See product label)

SPECIFICATION

Product Overview	Purified recombinant protein of full-length mouse CCAAT/enhancer binding protein (C/EBP), beta (Cebpb), with C-terminal MYC/DDK tag, was expressed in HEK293T cells.
Species	Mouse
Source	HEK293
Description	<p>Important transcription factor regulating the expression of genes involved in immune and inflammatory responses (PubMed:16585579, PubMed:17911624, PubMed:18486321, PubMed:20111005). Plays also a significant role in adipogenesis, as well as in the gluconeogenic pathway, liver regeneration, and hematopoiesis (PubMed:9727068, PubMed:10635333, PubMed:17301242, PubMed:17601773, PubMed:19478079, PubMed:24061474, PubMed:24216764). The consensus recognition site is 5'-T[TG]NNGNAA[TG]-3'. Its functional capacity is governed by protein interactions and post-translational protein modifications. During early embryogenesis, plays essential and redundant functions with CEBPA (PubMed:15509779). Has a prometotic effect on many cell types such as hepatocytes and adipocytes but has an antiproliferative effect on T-cells by repressing MYC expression, facilitating differentiation along the T-helper 2 lineage (PubMed:9727068, PubMed:10635333, PubMed:16585579). Binds to regulatory regions of several acute-phase and cytokines genes and plays a role in the regulation of acute-phase reaction and inflammation. Plays also a role in intracellular bacteria killing (PubMed:17911624). During adipogenesis, is rapidly expressed and, after activation</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

by phosphorylation, induces CEBPA and PPARG, which turn on the series of adipocyte genes that give rise to the adipocyte phenotype. The delayed transactivation of the CEBPA and PPARG genes by CEBPB appears necessary to allow mitotic clonal expansion and thereby progression of terminal differentiation (PubMed:15985551, PubMed:17301242, PubMed:17601773, PubMed:20194620). Essential for female reproduction because of a critical role in ovarian follicle development (PubMed:9303532). Restricts osteoclastogenesis (PubMed:19440205). Together with NFE2L1; represses expression of DSPP during odontoblast differentiation (By similarity).

Molecular Mass 31.9 kDa

Purity >80% as determined by SDS-PAGE and Coomassie blue staining

Stability Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

Storage Store at -80 centigrade after receiving vials.

Concentration >50 g/mL as determined by microplate BCA method

Storage Buffer 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.

GENE INFORMATION

Gene Name [Cebpb CCAAT/enhancer binding protein \(C/EBP\), beta \[Mus musculus \(house mouse\) \]](#)

Official Symbol [Cebpb](#)

Synonyms Cebpb; CCAAT/enhancer binding protein (C/EBP), beta; LAP; LIP; CRP2; NF-M;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Nfil6; NF-IL6; IL-6DBP; C/EBPbeta; CCAAT/enhancer-binding protein beta; AGP/EBP; C/EBP BETA; interleukin-6-dependent-binding protein; liver-enriched transcriptional activator; nuclear protein Il6

Gene ID [12608](#)

mRNA Refseq [NM_009883](#)

Protein Refseq [NP_034013](#)

UniProt ID [P28033](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA