

Cynomolgus monkey Cecum Lysate

Cat. No. Cecum-62C **Lot. No.** (See product label)

SPECIFICATION

Species	Cynomolgus
Source	Cynomolgus Cecum
Concentration	2 mg/mL or better
Tissue Type	Cecum
Shipping	Dry ice
Preparation method	<p>Monkey (Cynomolgus) cecum tissue lysate was prepared by homogenization using a proprietary technique. The tissue was frozen in liquid nitrogen immediately after excision and then stored at -70°C. The monkey (Cynomolgus) cecum tissue total protein is provided in a buffer including HEPES (pH 7.9), MgCl₂, KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, NP-40, and a cocktail of protease inhibitors. For quality control purposes, the cecum tissue pattern on SDS-PAGE gel is shown to be consistent for each lot by visualization with coomassie blue staining. The cecum tissue is then Western analyzed by either GAPDH or β-actin antibody, and the expression level is consistent with each lot.</p>
Diagnosis	Normal
Storage Buffer	HEPES pH 7.9, MgCl ₂ , KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, and NP-40.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Storage Instruction

Store at 2-8°C for continuous use. For extended storage, freeze working aliquots at -70°C. Repeated freezing and thawing is not recommended. Under proper storage conditions the shelf life is half a year from the date of receipt.

Applications

Cecum lysate is for use in Western blotting, 10 µg to 20 µg per lane is recommended for mini gel.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA