

Recombinant Mouse Chemokine (C-X3-C Motif) Ligand 1

Cat. No. Cx3cl1-127M **Lot. No.** (See product label)

SPECIFICATION

Species	Mouse
Source	Sf21 Cells
Description	<p>Fractalkine (FKN) is a transmembrane mucin-chemokine hybrid molecule expressed on activated endothelium that mediates attachment and firm adhesion of T cells, monocytes and NK cells. FKN plays an important proinflammatory role in rheumatoid arthritis (RA) pathogenesis as characterized by induction of synovial angiogenesis, chemotaxis, activation of monocytes and T cells as well as the stimulation of proliferation and synthesis of matrix degrading enzymes (matrix metalloproteinases, MMP) in synovial fibroblasts. Fractalkine thus may represent a novel target molecule for therapeutic intervention in RA.³ FKN is also an essential biomarker for predicting the prognosis of patients with colorectal cancer (CRC).</p>
State Of Matter	Lyophilized.
Purity	>97% by SDS Page and analyzed by silver stain.
Endotoxin	<1.0 EU/g as determined by the LAL method.
Molecular Weight	The predicted molecular weight of Recombinant Mouse Fractalkine is Mr 34 kDa. However, the actual molecular weight as observed by migration on SDS Page is Mr 90 kDa.
Storage And Stability	This lyophilized protein is stable for six to twelve months when stored desiccated at -

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

20°C to -70°C. After aseptic reconstitution, this protein may be stored at 2°C to 8°C for one month or at -20°C to -70°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.

GENE INFORMATION

Gene Name	Cx3cl1 chemokine (C-X3-C motif) ligand 1 [Mus musculus]
Synonyms	chemokine (C-X3-C motif) ligand 1; CX3C; Cxc3; Scyd1; ABCD-3; AB030188; A1848747; D8Bwg0439e; Cx3cl1; fractalkine; neurotactin; small inducible cytokine subfamily D, 1
Gene ID	20312
mRNA Refseq	NM_009142
Protein Refseq	NP_033168
UniProt ID	O35188
Chromosome Location	8 46.0 cM
Pathway	Chemokine signaling pathway; Cytokine-cytokine receptor interaction
Function	chemokine activity; cytokine activity; protein binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA