

Recombinant Human DDB2 Protein, His-tagged

Cat. No. DDB2-452H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human DDB2(Pro166~Met383) fused with His tag at N-terminal was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	Pro166~Met383
Description	This gene encodes a protein that is necessary for the repair of ultraviolet light-damaged DNA. This protein is the smaller subunit of a heterodimeric protein complex that participates in nucleotide excision repair, and this complex mediates the ubiquitylation of histones H3 and H4, which facilitates the cellular response to DNA damage. This subunit appears to be required for DNA binding. Mutations in this gene cause xeroderma pigmentosum complementation group E, a recessive disease that is characterized by an increased sensitivity to UV light and a high predisposition for skin cancer development, in some cases accompanied by neurological abnormalities. Two transcript variants encoding different isoforms have been found for this gene.
Form	PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Molecular Mass	28.6kDa
Identity	Reconstitute in PBS or others

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	<1.0EU per 1g (determined by the LAL method)
Purity	> 95%
Applications	Positive Control; Immunogen; SDS-PAGE; WB. If bio-activity of the protein is needed, please check active protein
Stability	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 centigrade for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Storage	Avoid repeated freeze/thaw cycles. Store at 2-8 centigrade for one month. Aliquot and store at -80 centigrade for 12 months.
Reconstitution	Reconstitute in PBS or others

GENE INFORMATION

Gene Name	DDB2 damage-specific DNA binding protein 2, 48kDa [Homo sapiens]
Official Symbol	DDB2
Synonyms	DDB2; damage-specific DNA binding protein 2, 48kDa; damage specific DNA binding protein 2 (48kD); DNA damage-binding protein 2; DDB p48 subunit; DDBB; FLJ34321; UV damaged DNA binding protein 2; UV DDB2; xeroderma pigmentosum group E protein; UV-DDB 2; UV-damaged DNA-binding protein 2; damage-specific DNA-binding protein 2; UV-DDB2;
Gene ID	1643

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq [NM_000107](#)

Protein Refseq [NP_000098](#)

MIM [600811](#)

UniProt ID [Q92466](#)



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA