Recombinant Human DDIT3 293 Cell Lysate

Cat. No. DDIT3-7027HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for DNA-damage-inducible transcript 3 (DDIT3) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	DDIT3 DNA-damage-inducible transcript 3 [Homo sapiens]
Official Symbol	DDIT3
Synonyms	DDIT3; DNA-damage-inducible transcript 3; DNA damage-inducible transcript 3 protein; C/EBP zeta; CHOP; CHOP10; GADD153; DDIT-3; c/EBP-homologous protein 10; CCAAT/enhancer-binding protein homologous protein; growth arrest and DNA damage-inducible protein GADD153; CEBPZ; CHOP-10; MGC4154;
Gene ID	1649
mRNA Refseq	NM_001195053
Protein Refseq	NP_001181982
MIM	126337
UniProt ID	P35638
Chromosome Location	12q13.1-q13.2
Pathway	ATF-2 transcription factor network, organism-specific biosystem; Activation of Chaperone Genes by ATF6-alpha, organism-specific biosystem; Activation of Chaperones by ATF6-alpha, organism-specific biosystem; Activation of Genes by ATF4, organism-specific biosystem; Adipogenesis, organism-specific biosystem; Diabetes pathways, organism-specific biosystem; Disease, organism-specific

	biosystem;
Function	DNA binding; protein dimerization activity; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; transcription corepressor activity; transcription factor binding;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA