

Recombinant Mouse EFNA2 cell lysate

Cat. No. EFNA2-2004MCL **Lot. No.** (See product label)

SPECIFICATION

| | |
|--------------------------------|---|
| Product Overview | Mouse EFNA2 / Ephrin-A2 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls |
| Species | Mouse |
| Source | Human Cells |
| Preparation method | Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized. |
| Lysis buffer | Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF |
| Quality control Testing | 12.5% SDS-PAGE Stained with Coomassie Blue |
| Recommended Usage | 1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[Efna2 ephrin A2 \[Mus musculus \]](#)

Official Symbol

EFNA2

Synonyms

EFNA2; ephrin A2; ephrin-A2; ephrin A6; CEK7-ligand; EPH related receptor tyrosine kinase ligand 6; EPH-related receptor tyrosine kinase ligand 6; Elf1; Epl6; CEK7L; Eplg6; Lerk6;

Gene ID

[13637](#)

mRNA Refseq

[NM_007909](#)

Protein Refseq

[NP_031935](#)

Pathway


Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem;

Function

ephrin receptor binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA