

Recombinant Human EGF, Fc-tagged

Cat. No. EGF-232H Lot. No. (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human EGF (aa 971-1023) is fused to the N-terminus of the Fc region of human IgG1.
Species	Human
Source	CHO
ProteinLength	971-1023 a.a.
Description	<p>Epidermal growth factor (EGF) is a growth factor and the founding member of the EGF family. All EGF family members are synthesized as type I transmembrane precursor proteins that may contain several EGF domains in the extracellular region. The mature proteins are released from the cell surface by regulated proteolysis. EGF is present in various body fluids, including blood, milk, urine, saliva, seminal fluid, pancreatic juice, cerebrospinal fluid, and amniotic fluid. Four ErbB (HER) family receptor tyrosine kinases including EGFR/ErbB1, ErbB2, ErbB3 and ErbB4, mediate responses to EGF family members. These receptors undergo a complex pattern of ligand induced homo or heterodimerization to transduce EGF family signals. EGF binds to the receptor EGFR stimulating the intrinsic protein-tyrosine kinase activity of the receptor. The tyrosine kinase activity initiates a signal transduction cascade that results in a variety of biochemical changes within the cell, including a rise in intracellular calcium levels, increased glycolysis and protein synthesis, and increases in the expression of certain genes including the gene for EGFR, which lead to DNA synthesis, cell growth, proliferation and differentiation. Other biological activities</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

ascribed to EGF include epithelial development, angiogenesis, inhibition of gastric acid secretion, fibroblast proliferation, and colony formation of epidermal cells in culture. Defects in EGF are the cause of hypomagnesemia type 4 (HOMG4), also known as renal hypomagnesemia normocalciuric. HOMG4 is a disorder characterized by massive renal hypomagnesemia and normal levels of serum calcium and calcium excretion. Clinical features include seizures, mild-to moderate psychomotor retardation, and brisk tendon reflexes.

Form	Lyophilized from 0.2µm-filtered solution in PBS.
Purity	≥98% (SDS-PAGE)
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Storage	Short Term Storage: +4°C; Long Term Storage: -20°C. Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name	EGF epidermal growth factor [Homo sapiens]
Official Symbol	EGF
Synonyms	EGF; epidermal growth factor; epidermal growth factor (beta urogastrone); pro-epidermal growth factor; beta-urogastrone; URG; HOMG4;
Gene ID	1950
mRNA Refseq	NM_001178130
Protein Refseq	NP_001171601

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	131530
UniProt ID	P01133
Chromosome Location	4q25
Pathway	Arf6 signaling events, organism-specific biosystem; Bladder cancer, organism-specific biosystem; Bladder cancer, conserved biosystem; Ceramide signaling pathway, organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Delta-Notch Signaling Pathway, organism-specific biosystem;
Function	calcium ion binding; epidermal growth factor receptor binding; growth factor activity; protein binding; transmembrane receptor protein tyrosine kinase activator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA