

Active Recombinant Human EIF2C2, His-tagged

Cat. No. EIF2C2-2180H Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant Human AGO2(Met 1-Ala 859), fused with His tag at N-terminal, was expressed in Baculovirus-Insect Cells.

Species Human

Source Insect Cells

ProteinLength Met 1-Ala 859

Description

Argonaute 2 (AGO2), also known as Eukaryotic translation initiation factor 2C2 (EIF2C2), belongs to the Argonaute family, AGO subfamily, which is a component of the RNA-induced silencing complex (RISC) and mediates small interfering RNA (siRNA)-directed mRNA cleavage and microRNA translational suppression. AGO2 protein is the catalytic engine of mammalian RNAi. It contains a PIWI domain that is structurally related to RNases H and possibly shares with them a two-metal-ion catalysis mechanism. Human AGO2 was unable to cleave preformed RNA duplexes and exhibited weaker binding affinity for RNA duplexes compared with the single strand RNA. The enzyme exhibited greater RNase H activity in the presence of Mn²⁺ compared with Mg²⁺. Human AGO2 exhibited weaker binding affinities and reduced cleavage activities for antisense RNAs with either a 5'-terminal hydroxyl or abasic nucleotide. In mouse hematopoiesis, AGO2 controls early development of lymphoid and erythroid cells. AGO2 is a highly specialized member of the Argonaute family with an essential nonredundant Slicer-independent function within the mammalian miRNA pathway. AGO2 regulates dFMR1 expression, and the relationship between dFMR1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

and AGO2 was defined by their physical interaction and co-regulation of downstream targets. AGO2 and dFMR1 are also connected through a regulatory relationship. AGO2 is a regulator of dFMR1 expression and have clarified an important developmental role for AGO2 in the nervous system and germ line that requires dFMR1 function. In addition, AGO2 is regulated at both the transcriptional and posttranslational level, and also implicate AGO2 and enhanced micro-RNA activity in the tumorigenic progression of breast cancer cell lines.

Predicted N Terminal Met

Form Lyophilized from sterile 20mM Tris, 500mM NaCl, pH7.4, 10% glycerol, 2mM DTT1. Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA.2. Please contact us for any concerns or special requirements.

Bio-activity Human AGO2 can bind Let-7a RNA and cleave target RNA (21nt).

Molecular Mass The recombinant human AGO2 consists of 877 amino acids and predicts a molecular mass of 99 kDa as estimated by SDS-PAGE under reducing conditions.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Purity > 85 % as determined by SDS-PAGE.

Stability Samples are stable for up to twelve months from date of receipt at -70 centigrade.

Storage Store it under sterile conditions at -20 centigrade to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	AGO2 argonaute RISC catalytic component 2 [Homo sapiens]
Official Symbol	AGO2
Synonyms	Q10; EIF2C2; protein argonaute-2; CTA-204B4.6; PAZ Piwi domain protein; PPD; argonaute 2; argonaute2; eIF-2C 2; eIF2C 2; eukaryotic translation initiation factor 2C, 2; hAgo2; protein slicer
Gene ID	27161
mRNA Refseq	NM_012154
Protein Refseq	NP_036286
MIM	606229
UniProt ID	Q9UKV8
Chromosome Location	8q24
Pathway	Adaptive Immune System, organism-specific biosystem; Ca ²⁺ pathway, organism-specific biosystem; Downstream signal transduction, organism-specific biosystem
Function	RNA 7-methylguanosine cap binding; RNA polymerase II core binding; core promoter binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA